

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 233/56, A61K 31/415, C07D 401/06, 403/06, 409/06, 405/06, 233/84, 413/12, 495/04 // (C07D 495/04, 335:00, 333:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/28300</p> <p>(43) International Publication Date: 10 June 1999 (10.06.99)</p>
<p>(21) International Application Number: PCT/US98/25669</p> <p>(22) International Filing Date: 3 December 1998 (03.12.98)</p> <p>(30) Priority Data: 08/985,347 4 December 1997 (04.12.97) US</p> <p>(71) Applicant: ALLERGAN SALES, INC. [US/US]; 2525 Dupont Drive, Irvine, CA 92612 (US).</p> <p>(72) Inventors: CHOW, Ken; 20 Tidal Surf, Newport Coast, CA 92657 (US). GIL, Daniel, W.; 2541 Point Del Mar, Corona Del Mar, CA 92625 (US). BURKE, James, A.; 670 West Third, Tustin, CA 92680 (US). HARCOURT, Dale, A.; 320 East Avenida, San Clemente, CA 92672 (US). GARST, Michael, E.; 2627 Raqueta, Newport Beach, CA 92660 (US). WHEELER, Larry, A.; 18 Valley View, Irvine, CA 92715 (US). MUNK, Stephen, A.; 15976 Winchester Drive, Northville, MI 48167 (US).</p> <p>(74) Agents: HOCH, James, M. et al.; Allergan Sales, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published</p> <p><i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>

(54) Title: SUBSTITUTED IMIDAZOLE DERIVATIVES HAVING AGONIST-LIKE ACTIVITY AT ALPHA 2B OR 2B/2C ADRENERGIC RECEPTORS



(57) Abstract

Compounds having adrenergic activity which are selective agonists for one or both of the α_{2B} and α_{2C} adrenoceptor receptor subtypes in preference to the α_{2A} adrenoceptor receptor subtype; the active compound being selected from the group consisting of compounds having formula (I) wherein the dotted lines represent optional bonds provided that two double bonds may not share a common carbon atom; R is H or lower alkyl; X is S or C(H)R¹, wherein R¹ is H or lower alkyl, but R¹ is absent when the bond between X and the ring represented by formula (a) is a double bond; Y is O, N, S, (CR²)₂, wherein y is an integer of from 1 to 3, -CH=CH- or -Y¹CH₂-, wherein Y¹ is O, N or S; x is an integer of 1 or 2, wherein x is 1 when R², R³ or R⁴ is bound to a saturated carbon atom and x is 2 when R², R³ or R⁴ is bound to a saturated carbon atom; R² is H, lower alkyl, halogen, hydroxy or lower alkoxy, or, when attached to a saturated carbon atom, R₂ may be oxo; R₃ and R₄ are, each, H, lower alkyl, hydroxy, lower alkoxy, or phenyl or, together, are -(C(R²)_x)_z-; -Y¹(C(R²)_x)_z-; -Y¹(C(R²)_x)_z-Y¹(C(R²)_x)_z-; -(C(R²)_x)_z-Y¹(C(R²)_x)_z- and -Y¹(C(R²)_x)_z-Y¹(C(R²)_x)_z- wherein z is an integer of from 3 to 5, z' is an integer of from 2 to 4 and x and y are as defined above, and further either end of each of these divalent moieties may attach at either R₃ or R₄ to form a condensed ring structure and the rings formed may be totally unsaturated, partially unsaturated, or totally saturated; and being useful for treating muscle spasticity including hyperactive micturition, diarrhea, diuretics, withdrawal syndromes, pain including neuropathic pain, neurodegenerative diseases, memory and cognition deficits, psychoses including manic disorders and anxiety, hypertension, cardiac ischemia, congestive heart failure, and nasal congestion without sedating or cardiovascular side effects.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**SUBSTITUTED IMIDAZOLE DERIVATIVES HAVING AGONIST-LIKE ACTIVITY AT ALPHA 2B
OR 2B/2C ADRENERGIC RECEPTORS**

5

1. Field of the Invention

The present invention is directed to a method of treating glaucoma or elevated intraocular pressure and other diseases with substantially reduced cardiovascular or sedative side effects by administering to
10 mammals including humans, compounds which are selective agonists of the $\alpha 2B$ alone or $\alpha 2B$ and $\alpha 2C$ adrenergic receptor subtypes and which lack substantial activity at the $\alpha 2A$ receptor subtype. The present invention is also directed to novel compounds and pharmaceutical compositions adapted for administering said compounds to mammals, including
15 humans.

2. Brief Description of the Prior Art

Compounds which have adrenergic activity are well known in the art, and are described in numerous United States and foreign patents and in scientific publications. It is generally known and accepted in the art that
20 adrenergic activity is useful for treating animals of the mammalian species, including humans, for curing or alleviating the symptoms and conditions of numerous diseases and conditions. In other words, it is generally accepted in the art that pharmaceutical compositions having an adrenergic compound or compounds as the active ingredient are useful for treating
25 glaucoma, chronic pain, nasal congestion, high blood pressure, congestive heart failure and inducing anesthesia.

The two main families of adrenergic receptor are termed alpha adrenergic receptors and beta adrenergic receptors in the art, and each of these two families is known to have subtypes, which are designated by

letters of the alphabet, such as $\alpha 2A$, $\alpha 2B$. See the article by Bylund et al, *Pharmacol Rev.* 46, pp. 121-136(1994).

SUMMARY OF THE INVENTION

5 It has been discovered in accordance with the present invention that adrenergic compounds which act selectively, and preferably even specifically as agonists of the $\alpha 2B$ or $\alpha 2B / \alpha 2C$ (hereinafter referred to as $\alpha 2B$ or $\alpha 2B/2C$) receptor subtypes in preference over the $\alpha 2A$ receptor subtype, possess desirable therapeutic properties associated with
10 adrenergics but without having one or more undesirable side effects such as changes in blood pressure or sedation. For the purposes of the present invention, a compound is defined to be a specific or at least selective agonist of the $\alpha 2B$ or $\alpha 2B/2C$ receptor subtype(s) if the compound is at least approximately ten times more potent as an agonist at either the $\alpha 2B$
15 and $\alpha 2C$ or both receptor subtypes than at the $\alpha 2A$ receptor subtype, or if the difference in the compound's efficacy at the $\alpha 2B$ and $\alpha 2B/2C$ receptor relative to the $\alpha 2A$ receptor is greater than 0.3 and its efficacy at the $\alpha 2A$ receptor is ≤ 0.4 .

Accordingly, the present invention relates to methods of treating
20 animals of the mammalian species, including humans, with a pharmaceutical composition comprising one or more specific or selective $\alpha 2B$ or $\alpha 2B/2C$ adrenergic agonist compounds as the active ingredient, for treatment of the many diseases or conditions against which alpha adrenergic compounds are useful, including without limitation glaucoma,
25 reducing elevated intraocular pressure, chronic pain, diarrhea, and nasal congestion. In addition, the compounds of this invention are useful for

treating muscle spasticity including hyperactive micturition, diarrhea, diuresis, withdrawal syndromes, pain including neuropathic pain, neurodegenerative diseases including optic neuropathy, spinal ischemia and stroke, memory and cognition deficits, attention deficit disorder,
5 psychoses including manic disorders, anxiety, depression, hypertension, congestive heart failure, cardiac ischemia and nasal congestion.

The present invention is also directed to the pharmaceutical compositions used in the above-noted methods of treatment.

The present invention particularly covers methods for treating
10 diseases and conditions where adrenergic compounds are effective for treatment, but their use is limited because of their generally known side effects.

DETAILED DESCRIPTION OF THE INVENTION

15 Compounds which are used in the pharmaceutical compositions and methods of treatment of the present invention are selective or specific agonists of the $\alpha 2B$ or $\alpha 2B/2C$ adrenergic receptor subtypes, in preference over the $\alpha 2A$ receptor subtype. In accordance with the present invention, a compound is considered a selective $\alpha 2B$ or $\alpha 2B/2C$ agonist if that
20 compound's difference in efficacy as an agonist of the $\alpha 2B$ or $\alpha 2B/2C$ receptor subtype(s) compared to the $\alpha 2A$ receptor subtype is greater than 0.3 and its efficacy at the $\alpha 2A$ receptor subtype is ≤ 0.4 and/or it is at least approximately 10 times more potent. Preferably, the compounds utilized in accordance with the present invention are specific agonists of the $\alpha 2B$ or
25 $\alpha 2B/2C$ receptor subtypes. Specifically, in this regard, a specific agonist is defined in the sense that a specific α adrenergic agonist does not

act as an agonist of the $\alpha 2A$ receptor subtype to any measurable or biologically significant extent.

A set of agents has been discovered that are functionally selective for the $\alpha 2B$ or $\alpha 2B/2C$ - subtypes of said adrenergic receptors. This preferential activity can be determined in a variety of functional assays such as Cyclic AMP Production, Shimizu et al, *J. Neurochem.* 16, pp. 1609-1619 (1969); R-SAT (Receptor Selection and Amplification Technology), Messier et al, *Pharmacol. Toxicol.* 76, pp. 308-311(1995) and the Cytosensor microphysiometer, Neve et al, *J. Biol. Chem.* 267, pp. 25748-25753, (1992) using cells that naturally express individual subtypes or have had one of the subtypes introduced. The cells or recombinant receptors used should be human or from a species that has been shown to have a similar pharmacology. In the study below, the RSAT assay on cells that have been transiently transfected with the human $\alpha 2A$ (c10 gene), rat $\alpha 2B$ (RNG gene) and human $\alpha 2C$ (c4 gene) receptors was used. The rat $\alpha 2B$ receptor has been shown to have a pharmacology that corresponds to the human $\alpha 2B$ receptor (see, for example, Bylund et al., *Pharmacol. Rev.* 46, pp. 127-129(1994)).

In the treatment of glaucoma, particularly, topical administration may be used. Any common topical formulation such as a solution, suspension, gel, ointment, or salve and the like may be applied to the eye in glaucoma and dermally to treat other indications. Preparation of such topical formulations are well described in the art of pharmaceutical formulations as exemplified, for example, by Remington's Pharmaceutical Science, Edition 17, Mack Publishing Company, Easton, Pennsylvania.

If the drug is to be administered systemically, it may be confectioned as a powder, pill, tablet or the like or as a syrup or elixir for oral administration. For intravenous, intraperitoneal, intrathecal or epidural administration, the compound will be prepared as a solution or suspension capable of being administered by injection. In certain cases, it may be useful to formulate these compounds in suppository or as an extended release formulation, including the dermal patch form, for deposit on or under the skin or for intramuscular injection.


Treatment of glaucoma or any other indications known or discovered to be susceptible to treatment by adrenergic compounds will be effected by administration of therapeutically effective dose of one or more compounds in accordance with the instant invention. A therapeutic concentration will be that concentration which effects reduction of the particular condition, or retards its expansion. In certain instances, the drug potentially could be used in a prophylactic manner to prevent onset of a particular condition. A given therapeutic concentration will vary from condition to condition and in certain instances may vary with the severity of the condition being treated and the patient's susceptibility to treatment. Accordingly, a given therapeutic concentration will be best determined at the time and place through routine experimentation. However, it is anticipated that in the treatment of, for example, glaucoma, that a formulation containing between 0.001 and 5 percent by weight, preferably about 0.01 to 3% will usually constitute a therapeutically effective concentration. If administered systemically, an amount between 0.001 and 50 mg per kg, preferably between 0.001 and 10 mg per kg body weight per

day, but most preferably about 0.01 to 1.0 mg/kg, will effect a therapeutic result in most instances.

Because the $\alpha 2B$ and $\alpha 2B/2C$ specific selective agonist compounds lack substantial $\alpha 2A$ side effects, treatments of diseases or conditions with such compounds in accordance with the present invention is advantageous, particularly when the treatment is directed to a human having cardiovascular problems.

The general structures of exemplary specific $\alpha 2B$ and $\alpha 2C$ agonist or selective $\alpha 2B$ and $\alpha 2B/2C$ agonist adrenergic compounds which are used in the pharmaceutical compositions and methods of treatment of the present invention are provided by general Formulas, below.

In one aspect of the invention, a compound having selective agonist activity at the $\alpha 2B$ or $\alpha 2B/2C$ adrenergic receptor subtype(s) as compared to the $2A$ adrenergic receptor subtype is represented by the general formula



I


wherein the dotted lines represent optional bonds provided that two double bonds may not share a common carbon atom; R is H or lower alkyl; X is S or C(H)R', wherein R' is H or lower alkyl, but R' is absent when the bond between X and the ring represented by

7



- is a double bond; Y is O, N, S, $(CR^1)_y$, wherein y is an integer of from 1 to 3,
 $-CH=CH-$ or $-Y^1CH_2-$, wherein Y^1 is O, N or S; x is an integer of 1 or 2,
 wherein x is 1 when R^2 , R^3 or R^4 is bound to an unsaturated carbon atom
 5 and x is 2 when R^2 , R^3 or R^4 is bonded to a saturated carbon atom; R^2 is H,
 lower alkyl, halogen, hydroxy or lower alkoxy, or, when attached to a
 saturated carbon atom, R_2 may be oxo; R_3 and R_4 are, each, H, lower alkyl,
 hydroxy, lower alkoxy, or phenyl or, together, are $-(C(R^2)_x)_z-$; -
 $Y^1(C(R^2)_x)_z-$; - $Y^1(C(R^2)_x)_y Y^1-$; $-(C(R^2)_x)- Y^1-(C(R^2)_x)-$; $-(C(R^2)_x)- Y^1-$
 10 $(C(R^2)_x)-(C(R^2)_x)-$ and $- Y^1-(C(R^2)_x)- Y^1-(C(R^2)_x)-$ wherein z is an integer of
 from 3 to 5, z' is an integer of from 2 to 4 and x and y are as defined above,
 and further either end of each of these divalent moieties may attach at
 either R^3 or R^4 to form a condensed ring structure shown generally as


15



- and the rings formed may be totally unsaturated, partially unsaturated, or
 20 totally saturated provided that a ring carbon has no more than 4 valences,
 nitrogen no more than three and O and S have no more than two.

In another aspect of the invention in the above compound is
 represented by the formula

25



wherein X may be C(H)R' and R' is H.


In said compound of formula II, R₂ may be H and



may represent a furanyl radical.

In such furanyl derivatives of Formula II, R³ and R⁴ together may be (CH)₂, or R³ may be H and R⁴ may be t-butyl, or R³ and R⁴ may be H, or R³ may be H and R⁴ may be methyl or ethyl.

10 Alternatively, in the compound of Formula I, R¹ may be methyl and



may represent a furanyl radical.

15 Alternatively, in said compounds of Formula II, R² may be H and



may represent a thienyl radical.

20 In such thienyl derivatives of Formula II, R³ and R⁴, together, may represent (CH)₂, or R³ may be phenyl and R⁴ may be H, or R³ and R⁴, together, may represent (CH)₂S, or R³ and R⁴ may be H, or R³ and R⁴, together, may represent (CH)₂, or may be R³ may be H and R⁴ may be methyl, or R³ may be bromo and R⁴ may be H, or R³ may be hydrogen and

25 R⁴ may be chloro, or R³ may be methyl and R⁴ may be hydrogen.

Alternatively, in the compounds of Formula II

9



may represent a cyclohexyl radical.

- In such cyclohexyl derivatives of Formula II, R^2 may be hydrogen and R^3 and R^4 may, together, represent $(CH)_n$, or R^2 may be oxo and R^3 and R^4 , together, may be $(CH)_n$, or R^2 may be hydrogen or oxo and R^3 and R^4 , together, may represent $(CH)_nS$, or R^2 may be hydrogen and R^3 and R^4 may, together, represent $(CH)_n$, forming an octahydronaphthalene, or R^2 may be oxo and R^3 and R^4 may, together, represent $(CH)_n$, or R^2 may be oxo and R^3 and R^4 , together, may represent $(CH)_n C(CH_3)(CH)$, or R^2 may be hydrogen and R^3 and R^4 , together, may represent $S(CH_2)_n$, or R^2 , R^3 and R^4 may be H, or R^2 may be oxo and R^3 and R^4 , together, may represent $(CH)_2 C(OCH_3)CH$, or R^3 and R^4 together may represent $-Y^1-C(R_2)_n-C(R_2)_n-$ wherein Y^1 is N, forming a tetrahydroquinoxaline wherein R^2 may be hydrogen or oxo.
- Alternatively, in the compounds of Formula II

15



- may represent a tetrahydroquinoline radical wherein R^3 and R^4 together are $-Y^1-C(R_2)_n-C(R_2)_n-C(R_2)_n-$ wherein Y^1 is N. In such tetrahydroquinoline derivatives $(R^2)_n$ may be hydrogen or oxo; or may represent a tetrahydro-isoquinoline radical wherein R^3 and R^4 together are $-C(R_2)_n-Y^1-C(R_2)_n-C(R_2)_n-$ wherein Y^1 is N and $(R^2)_n$ may be hydrogen or oxo.

Alternatively, in the compounds of Formula II




may represent a cyclopentyl radical.

In such cyclopentyl derivatives of Formula II, R^2 may be H and R^3 and R^4 , together, may represent $(CH)_n$, or R^2 may be oxo and R^3 and R^4 , together, may represent $(CH)_n$, or R^2 may be hydrogen and R^3 and R^4 , together, may represent $(CH)_n$.

In another aspect of the invention, Y is $(CH)_n$, and X may be CH and R^2 may be oxo or X may be CH_2 and R^2 may be H. Alternatively, R^3 and R^4 , together, may represent $(CH)_n$, Y may be $CH_2C(CR^1)_n$, wherein R^1 is hydrogen, or Y may be $-CH_2C(Me)-$ and R^2 may be hydrogen or oxo.


Finally, in the compounds of Formula II



may represent a phenyl radical.

In such phenyl derivatives of Formula I, X may be CH_2 , R may be H or CH_3 , R^2 , R^3 and R^4 may be H, or R^3 and R^4 , together, represent $O(CR^1)_nO$ to provide a 1,4-benzodioxan derivative, or alternatively, X may be S and R^2 , R^3 and R^4 may be H.

In another aspect of the invention, said compound has the formula




wherein Y is S or O.

In such compound of Formula III, X may be C(H)R¹, R, R¹, R², R³ and R⁴ may be H and Y may be O or S.

In another aspect of the invention, said compound has the formula

5

10



and R³ and R⁴, together, represent (CH)₂.

15


In such compounds of Formula IV, Y¹ may be O, R² may be oxo and X is CH or CH₂, or one of R³ is hydroxy and the other may be H, or R² may be H.

In such compounds of Formula IV, Y¹ may be S, X may be CH₂ and R² may be oxo, or R² may be H and X may be CH and R² may be oxo.

20

In another aspect of the invention, the compound having selective activity at the 2B or 2B and 2C adrenergic receptor subtype(s) as compared to the 2A adrenergic receptor subtype is represented by the formula


25




alternatively W is a bicyclic radical selected from the group consisting of

12

5




- wherein R^5 , R^6 , R^7 and R^8 are selected from the group consisting of H and
 10 lower alkyl provided that at least one of R^5 and R^6 or R^7 and R^8
 are $OC(R^9)C(R^9)N(R)$ to form a condensed ring with




wherein R^9 is H, lower alkyl or oxo;
 and

15




- 20 wherein R^{10} is H, lower alkyl, phenyl or lower alkyl substituted phenyl,
 and Z is O or NH. Compounds wherein W is norbornyl are disclosed and
 claimed in commonly assigned co-pending application 09/003902, filed on
 7 January, 1998, which is hereby incorporated by reference in its entirety.

In one aspect of the invention Z may be O and W may be




and R^{10} may be selected from the group consisting of H, phenyl and o-methylphenyl, e.g. R^{10} may be o-methylphenyl.

In another aspect of the invention W may be



- 5 wherein Z may be NR, R may be methyl or hydrogen, one of $(R^9)_x$ may be H and R^5 may be H.

Alternatively, W may be




wherein R may be H and R^8 may be methyl.

- 10 The invention is further illustrated by the following examples (including general synthetic schemes therefore) which are illustrative of

various aspects of the invention and are not intended as limiting the scope of the invention as defined by the appended claims.

Example A

- 5 Synthesis of 1-dimethylsulfamoyl-2-t-butyltrimethylsilyl-5-imidazolecarboxaldehyde:



Procedure -


- 10 Imidazole (1) (20.0g, 0.29 mol), triethylamine (41.0mL, 0.29 mol) and N,N-dimethylsulfamoyl chloride (31.6mL, 0.29 mol) were added to 320mL of benzene. The reaction was stirred for 48h at room temperature (rt) and then filtered. The filtrate was collected and concentrated under reduced pressure. Vacuum distillation of the crude product (~ 0.5 mmHg, 115° -
 15 118°C) afforded 38.7g (76%) of a clear and colorless oil. Upon cooling the product solidifies to give white crystals (2). 1-(Dimethylsulfamoyl)imidazole (2) (18.8g, 0.11 mol) was added to 430mL of tetrahydrofuran (THF). The solution was cooled to -78°C . A solution of n-butyl lithium (n-BuLi) in hexane (1.6M, 70.9 mL, 0.11 mol) was added dropwise to the

reaction flask. Upon completion, the reaction was stirred for 1h at -78°C. t-Butyldimethylsilylchloride (17.8g, 0.12 mol) in 50mL of THF was added via cannula to the reaction. After the addition was completed the reaction mixture was warmed slowly to rt and then stirred for 24h. The reaction was diluted with water and the organic layer separated. The organic phase was washed with brine and then dried over sodium sulfate. The mixture was filtered and the filtrate concentrated under reduced pressure. Column chromatography (20% ethyl acetate/ hexane as eluant) afforded a light yellow solid. Recrystallization from pentane gave 30g (94%) of white crystals (3).

1-Dimethylsulfamoyl-2-t-butyldimethylsilyl imidazole (3) (5.0g, 17.3 mmol) was added to 100mL of THF. The solution was cooled to -20°C. A solution of secondary butyl lithium (s-BuLi) in hexane (1.3M, 14.6mL, 19 mmol) was added dropwise to the reaction flask. Upon completion the reaction was stirred for 1h at -20°C. 8 mL of dimethylformamide (DMF) was added to the reaction and then stirred at rt for 3.5h. The reaction was diluted with water and the organic layer separated. The organic phase was washed with brine and then dried over sodium sulfate. The mixture was filtered and the filtrate concentrated under reduced pressure. Column chromatography (20% ethyl acetate/ hexane) afforded a light yellow oil. Upon cooling the product solidifies to give yellow crystals of 1-dimethylsulfamoyl-2-t-butyldimethylsilyl-5- imidazolecarboxaldehyde (4).

Example B-1

Procedure for Preparation of 4(5)-(7-methoxy-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole, hydrogen chloride salt:



5

Procedure -

7-Methoxy-1-tetralone (1) (1.5g, 8.5 mmol) and 1-(dimethylsulfamoyl)-2-(tert-butyldimethylsilyl)-5-imidazolecarboxaldehyde (2) (2.7g, 8.5 mmol) were added to 8.5 mL of a 40% solution of sulfuric acid. The reaction was heated for 24h at 90°C. After cooling to rt, the reaction was made basic with excess concentrated ammonium hydroxide. The mixture was extracted twice with THF. The organic layers were combined and washed

10

with brine. The organic layer was separated and dried over sodium sulfate. The mixture was filtered and the filtrate concentrated under reduced pressure to afford 2.7g of a yellow solid (3) comprising 3-(3H-imidazole-4(5)ylmethylene)-7-methoxy chroman-4-one. The crude product
5 was suspended in 100mL of ethanol and a palladium on carbon catalyst (10%, 0.27g) added. The mixture was shaken in a Parr hydrogenator apparatus while under 40 psi of hydrogen. After 19h the reaction mixture was filtered through Celite and the filtrate concentrated under reduced pressure. Column chromatography with 7% methanol in chloroform
10 afforded 1.05g (46%) of a tan color solid comprising 2-[3H-Imidazole-4(5)-ylmethyl]-7-methoxy-3,4-dihydro-2H-naphthalen-1-one (4)(B-1a). (4) (0.5g, 1.95 mmol) was added to 20mL of methanol. Sodium borohydride (74mg, 1.95 mmol) was added to the solution. After stirring for 2.5h at rt the reaction mixture was quenched with water. The reaction mixture was then
15 extracted twice with ethyl acetate. The organic layers were combined and washed with brine. The organic layer was separated and dried over sodium sulfate. The mixture was filtered and the filtrate concentrated under reduced pressure to afford 0.5g of a white solid (5) comprising 2-[3H-Imidazole-4(5)-ylmethyl]-7-methoxy-3,4-dihydro-2H-naphthalen-1-ol.
20 The crude product was dissolved in 26mL of dichloromethane. Triethylsilane (2.5mL, 15.6 mmol) and trifluoroacetic acid (4.8mL, 62.3 mmol) were added and the reaction stirred at rt for 22h. The reaction was made basic with 2N NaOH and the organic layer separated and washed with brine. The solution was dried over sodium sulfate. The mixture was
25 filtered and the filtrate concentrated under reduced pressure. Column chromatography with 7% methanol in chloroform afforded 0.39g (83%) of a

tan color oil (6). The product was dissolved in methanol and an excess of hydrogen chloride (HCl) in ether was added. The solution was concentrated under reduced pressure to yield 0.3g of a tan color solid. Column chromatography with 7% methanol in chloroform afforded 0.25g (46%) of 4(5)-(7-methoxy-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole, hydrogen chloride salt (B-1) as white crystals (7) after recrystallization from a mixture of acetone and methanol.

¹H NMR (300 MHz, CD₃OD) 8.83 (s, 1H), 7.38 (s, 1H), 6.95 (d, 1H, J=8.5Hz), 6.66 (d, 1H, J=8.4Hz), 6.57 (s, 1H), 3.73 (s, 3H), 2.71-2.81 (m, 5H), 2.43-2.52 (m, 1H), 1.90-2.14 (m, 2H), 1.40-1.51 (m, 1H).

Following the procedure of Example B-1 various fused ring compounds are reacted to yield the imidazole derivatives listed below.

15		Example B-2(a-d)
	4-chromanone	(2a) 3-(3H-imidazol-4(5)-ylmethylene)chroman-4-one
20		(2b) 3-(3H-imidazol-4(5)-ylmethyl)chroman-4-one
		(2c) 3-(3H-imidazol-4(5)-ylmethyl)chroman-4-ol
25		(2d) 4(5)-chroman-3-ylmethyl-1H-imidazole
		Example B-3(a-b)
	1-tetralone	(3a) 2-(3H-imidazol-4(5)-ylmethyl)-3,4-dihydro-2H-naphthalen-1-one
30		(3b) 4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole

Example B-4(a-b)

- 5 4-methyl-1-tetralone (4a) 4(5)-(4-methyl-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole
- 10 (4b) 2-(3H-imidazol-4(5)-ylmethyl)-4-methyl-3,4-dihydro-2H-naphthalen-1-one

Example B-5(a-b)

- 15 Thiochroman (5a) 3-(3H-imidazol-4(5)-ylmethylene)thiochroman-4-one
- 20 (5b) 3-(3H-imidazol-4(5)-ylmethyl)thiochroman-4-one

Example B-6

25 The hydrogen chloride salt of the previous compound is prepared by step 5 of the method of Example B-1, above.

- 30 Thiochroman 4(5)-thiochroman-3-ylmethyl-1H-imidazole

Example B-7(a-c)

- 35 1-indanone (7a) 2-(3H-imidazol-4(5)-ylmethylene)indan-1-one
- 40 (7b) 2-(3H-imidazole-4(5)-ylmethyl)indan-1-one
- 45 (7c) 4(5)-indan-2-ylmethyl-1H-imidazole

Example B-8(a-b)


- 7-methyl-1-tetralone (8a) 2-(3H-imidazol-4(5)-ylmethyl)-7-methyl-3,4-dihydro-2H-naphthalen-1-one
- 5 (8b) 4(5)-(7-methyl-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole
- 10 The hydrogen chloride salt of this compound is prepared by the method of Example B-6.
- Example B-9(a-c)
- 15 4-keto-4,5,6,7-tetrahydrothianaphthene (9a) 4(5)-(4,5,6,7-tetrahydrobenzo[b]thiophen-5-ylmethyl)-1H-imidazole
- The hydrogen chloride salt of this compound is prepared by the method of Example B-6.
- 20 (9b) 5-(3H-imidazol-4(5)-ylmethyl)-6,7-dihydro-5H-benzo[b]thiophen-4-one
- The hydrogen chloride salt of this compound is prepared by the method of Example B-6.
- 25 (9c) 5-(octahydrobenzo[b]thiophen-5-ylmethyl)-1H-imidazole
- Example B-10
- 30 4,4-Dimethyl-1-tetralone 4(5)-(4,4-dimethyl-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole
- Example B-11(a-b)
- 35 1-Benzosuberone (11a) 4(5)-(6,7,8,9-tetrahydro-5H-benzocyclohepten-6-ylmethyl)-1H-imidazole

(11b) 6-(1H-imidazol-4(5)-ylmethylene)-6,7,8,9-tetrahydrobenzocyclohepten-5-one

5

Example C-1

Procedure for Preparation of 4(5)-thiophen-3-ylmethyl-1H-imidazole :



Procedure -

1-(Dimethylsulfamoyl)imidazole (1) (2.0g, 11.4 mmol) is taken up in 42mL of anhydrous THF and cooled to -78°C . $n\text{-BuLi}$ (6.6mL, 10.6 mmol) is added dropwise to the solution of (1). The resultant solution is stirred at -78°C for 30 min. Tert-butyldimethylsilylchloride (TBSCl) (1.6g, 10.6 mmol) in 8mL of THF is added to the reaction. The reaction is warmed to rt and stirred overnight. The next day the reaction is cooled to -20°C and 7.3mL

- (11.6 mmol) of *n*-BuLi added. After stirring at -20°C for 45 min, 3-thiophene carboxaldehyde (2) (1.0mL, 11.6 mmol) is added to the reaction mixture. Then reaction is warmed to rt and stirred overnight. The next day the reaction is quenched with water and diluted with ethyl acetate.
- 5 The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (2:5 ethyl acetate/ hexane) affords 3.0g (7.5 mmol) of 2-(*t*-butyldimethylsilyl)-5-(hydroxythiophen-2-ylmethyl)imidazole-1-sulfonic acid dimethylamide (3). (3) (1.5g, 3.74
- 10 mmol) is taken up in 37mL of THF. A 1M solution of tetra-*n*-butylammonium fluoride (TBAF) in THF (4.1mL, 4.1 mmol) is added dropwise to the solution of (3). The reaction is stirred overnight at rt. The next day the reaction is quenched with water and then extracted with ethyl acetate. The organic layer is washed with water followed by brine. The
- 15 organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. 0.94g (3.3 mmol) of 5-(hydroxythiophen-2-ylmethyl)imidazole-1-sulfonic acid dimethylamide (4) is recovered. (4) (0.5g, 1.74 mmol) is taken up in 23mL of dichloromethane, to the solution is added 2.2 mL (13.9 mmol) of triethylsilane and 4.3 mL (55.7 mmol) of
- 20 trifluoroacetic acid. The reaction is stirred at rt overnight and then quenched with water and neutralized with solid sodium bicarbonate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography using a 1:1 mixture of ethyl acetate and
- 25 hexane affords 0.42g (1.55 mmol) of 5-(thiophen-2-ylmethyl)imidazole-1-sulfonic acid dimethylamide (5). (5) (0.42g, 1.55 mmol) is taken up in 10mL

of a 1.5N HCl solution and heated at reflux for 3h and then stirred at rt overnight. The reaction is diluted with ethyl acetate, neutralized with solid sodium bicarbonate and then made basic with 2N NaOH. The organic layer is washed with water followed by brine. The organic phase is dried
5 over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography using a 10:1 mixture of chloroform and methanol affords 0.17g (1.0 mmol) of 4(5)-thiophen-3-ylmethyl-1H-imidazole (6) (C-1).

¹H NMR (300 MHz, CD₃OD) 7.52 (s, 1H), 7.25-7.27 (m, 1H), 6.96-7.01 (m,
10 2H), 6.77 (s, 1H), 3.98 (s, 2H).

Example C-2

The 2-carboxaldehyde isomer of 3-thiophene carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-thiophen-2-ylmethyl-1H-imidazole

15 Example C-3

5-Methyl-2-thiophene carboxaldehyde of 3-thiophene carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-(5-methylthiophen-2-ylmethyl)-1H-imidazole

Example C-4

20 5-Chloro-2-thiophene carboxaldehyde of 3-thiophene carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-(5-chlorothiophen-2-ylmethyl)-1H-imidazole

Example C-5

2-Furan carboxaldehyde is substituted into the method of Example C-1 to
25 yield 4 (5)-furan-2-ylmethyl-1H-imidazole

Example C-6

3-Furan carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-furan-3-ylmethyl-1H-imidazole

Example C-7

- 5 5-Methyl-2-furan carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-(5-methylfuran-2-ylmethyl)-1H-imidazole

Example C-8

Benzaldehyde is substituted into the method of Example C-1 to yield 4(5)-benzyl-1H-imidazole

10

Example C-9

2-Thianaphthene carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-benzo[b]thiophen-2-ylmethyl-1H-imidazole

Example C-10

- 15 2-Benzofuran carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-benzofuran-2-ylmethyl-1H-imidazole

Example C-11

5-Ethyl-2-furan carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-(5-ethylfuran-2-ylmethyl)-1H-imidazole

Example C-12

- 20 4-Bromo-2-thiophene carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-(4-bromothiophen-2-ylmethyl)-1H-imidazole

Example C-13

4-Phenyl-2-thiophene carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-(4-phenylthiophen-2-ylmethyl)-1H-imidazole

25


Example C-14

4-Methyl-2-thiophene carboxaldehyde is substituted into the method of Example C-1 to yield 4(5)-(4-methylthiophen-2-ylmethyl)-1H-imidazole,hydrochloride salt

5

Example D-1

Procedure for Preparation of oxazolidin-2-ylidene-(3-phenyl bicyclo[2.2.1]hept-2-yl) amine :

**10 Procedure -**

The *endo exo* relative stereochemistry of the compound was prepared, by making the β -nitrostyrene as shown above. Treatment of a methanol solution of benzaldehyde (10g, 94.3 mmole) with nitromethane (51ml, 943 mmol) in the presence of sodium hydroxide (3N in methanol to pH=8) afforded the nitro alcohol in 60% yield. Dehydration of the alcohol was effected by treatment with methanesulfonyl chloride (3.56g, 31.1mmole) followed by triethylamine (6.3g, 62.2 mmol) in

dichloromethane (35ml) to give 97% yield of product. Kugelrohr distillation was done to purify compound. Construction of the bicyclo[2.2.1]heptane skeleton was carried out in one step. The Diels-Alder reaction was conducted by warming the nitrostyrene (4.5g, 30.2 mmole) with cyclopentadiene (3.98g, 60.4 mmole) in 1, 2-dichloroethane (10ml). The Diels-Alder reaction proceeds in approximately a 3:1 endo:exo nitro ratio. Both the ratio and relative stereochemistry was demonstrated through x-ray analysis. Reduction of both the nitro group and the olefin was carried out under an atmosphere of hydrogen in the presence of 10% by weight palladium on charcoal. Separation of isomers was conveniently carried out at this stage using flash chromatography with 5% ammonia-saturated methanol in dichloromethane. The amine (0.7g, 3.74 mmole) was treated first with chloroethylisocyanate (0.38ml, 4.49mmole) to afford the chloroethylurea, which was then warmed in the presence of aqueous NaHCO₃ solution to afford oxazolidin-2-ylidene-(3-phenyl bicyclo[2.2.1]hept-2-yl) amine (D-1) in 51% yield.

¹H NMR (300 MHz, CDCl₃) δ 1.36-1.80 (m, 6H), 2.14 (d, 1H, J=4.40Hz), 2.37 (s, 1H), 2.65 (s, 1H), 3.71-3.78 (m, 2H), 3.95-3.98 (m, 1H), 4.19-4.25 (t, 2H, J=17.15Hz, J=8.36Hz), 7.17-7.29 (m, 5H).

20

Example D-2

Oxazolidin-2-ylidene-(3-o-tolyl bicyclo[2.2.1]hept-2-yl)amine is prepared by substituting o-methyl β-nitrostyrene in the method of D-1


Example D-3

Bicyclo[2.2.1]hept-2-yl oxazolidin-2-ylidene amine is prepared by substituting nitroethene in the method of D-1

25

Example E-1

Procedure for Preparation of imidazolidin-2-ylidene-(4-methyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)amine:



5

Procedure -

To 2-amino-4-nitrophenol (**1**) (4.00g, 25.95 mmol), triethylamine (15.20mL, 109.0 mmol) and 4-dimethylaminopyridine (0.063g, 0.52 mmol) slurried in anhydrous CH_2Cl_2 (250mL) at 0°C under argon added


chloroacetyl chloride (2.27 mL, 28.55mmol) via syringe. After refluxing for 72h pure product was filtered off and washed with water. The mother liquor was washed successively with phosphoric acid (0.5M), saturated sodium bicarbonate, water and brine and then dried over MgSO_4 . This solution was adhered to silica and purified by flash chromatography on silica with hexane/ethyl acetate (4:6) to give additional product. The combined solids were dried in vacuo to give pure 6-nitro-4H-benzo[1,4]oxazin-3-one (2) (4.12g) in 82% yield. To a slurry of (2) (1.49g, 7.65 mmol) in anhydrous THF (40mL) under argon in a 2-neck round-bottom flask equipped with a reflux condenser was added borane-dimethyl sulfide complex (15.3mL, 30.62 mmol). The mixture was heated at reflux until starting material was no longer observed via thin layer chromatography (2h). The reaction mixture was cooled to rt and carefully quenched by the dropwise addition of methanol. The resulting mixture was then refluxed an additional 10 minutes. The crude reaction mixture was concentrated in vacuo and purified by flash chromatography on silica with hexane/ethyl acetate (8:2) to give pure 6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine (3) (1.36g) as an orange solid in 99% yield. To (3) (0.032g, 0.178mmol) and formalin (37% in H_2O , 0.20 mL, 2.67 mmol) in anhydrous acetonitrile (1.5mL) at ambient temperature was added sodium cyanoborohydride (0.034g, 0.534 mmol). This solution was stirred for 30 min before adding glacial acetic acid (0.032mL, 0.534 mmol). The resulting mixture was stirred an additional 16h. The organics were taken up in diethyl ether and washed successively with NaOH (2N) and brine, dried over MgSO_4 and concentrated in vacuo. The resulting solids were purified by flash chromatography on silica with hexane/ethyl acetate (7:3) to give

pure 4-methyl-6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine (4) (0.031g) in 93% yield. To (4) (2.16g, 11.12 mmol) and 10% palladium on carbon (0.216g, 10 wt. %) under argon was added methanol (MeOH) (30mL) followed by THF (30mL). Hydrogen was bubbled thru the resulting slurry until no (4) remained visible by thin layer chromatography (2h). Celite was added and the mixture was filtered through a bed of celite followed by a MeOH wash. The resulting solution was concentrated in vacuo to give pure 4-methyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-ylamine (5) (1.86g) as a pale purple oil in 100% yield which was carried on without further purification. To (5) (1.86g, 11.34 mmol) and imidazoline-2-sulfonic acid (1.84g, 12.24 mmol) in anhydrous acetonitrile (50mL) under argon at 0°C was added triethylamine (3.26mL, 23.36 mmol). This solution was gradually warmed to ambient temperature and stirred for 16h. At that time an additional amount of imidazoline-2-sulfonic acid (0.86g, 5.55 mmol) was added and the resulting mixture was stirred an additional 5h. This solution was concentrated in vacuo and the residues were taken up in H₂O. The organics were extracted into CH₂Cl₂ and washed twice with NaOH and then brine, dried over MgSO₄ and concentrated in vacuo. The resulting foam was purified by flash chromatography on silica with 20% methanol (saturated with ammonia) in chloroform to give pure imidazolidin-2-ylidene-(4-methyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl)amine (6) (E-1) (0.905g) in 34% yield.

¹H NMR (CDCl₃): 2.81 (s, 3H); 3.26 (t, J=8.9 Hz, 2H); 3.60 (s, 4H); 4.26 (m, 2H); 4.60 (vbrs, 2H); 6.34 (dd, J=8.2 Hz, J=2.4 Hz, 1H); 6.39 (d, J=2.4 Hz, 1H); 6.68 (d, J= 8.2 Hz, 1H).

Example F & G

Procedure for Preparation of 6-(imidazolidin-2-ylidene amino)-5-methyl-4H-benzo[1,4]oxazin-3-one (F) and Imidazolidin-2-ylidene-(5-methyl-3,4-dihydro-2H-benzo[1,4]oxazin-8-yl)amine (G):



Procedure -

- To 2-amino-3-methylphenol (1) (14.72g, 0.120 mol), triethylamine (35.0mL, 0.251 mol) and 4-dimethylaminopyridine (0.29g, 2.39 mmol) in anhydrous CH_2Cl_2 (100mL) at 0°C under argon was added chloroacetyl chloride (10.0mL, 0.126 mol) dropwise via syringe. After the addition was complete the resulting solution was refluxed for 24h. The organics were washed successively with phosphoric acid (0.5M), saturated sodium bicarbonate, water and brine and then dried over MgSO_4 . The resulting solution was concentrated and taken up in THF to which ether was added.
- 10 The resulting crystals were filtered off to give pure 5-methyl-4H-benzo[1,4]oxazin-3-one (2) (12.30g) in 63% yield. To (2) (14.64g, 89.72 mmol) dissolved in concentrated H_2SO_4 (65 mL) at -10°C was added 70% concentrated HNO_3 (8.08g, 89.72 mmol) in concentrated H_2SO_4 (25mL) with rapid mechanical stirring at a rate whereby the internal temperature was
- 15 maintained below -5°C . As soon as the addition was complete the mixture was poured onto crushed ice (500mL) and the resultant solids were filtered off and slurried in cold water (300 mL) while sufficient NaOH was added to adjust the pH to 7. The recovered yellow powder was dissolved in THF, adhered to silica and purified by flash chromatography with 60% hexane
- 20 and ethyl acetate to give the nitrated product as a mixture of two regioisomers, i.e. the desired 6-substituted aromatic comprising 6-nitro-5-methyl-4H-benzo[1,4]oxazin-3-one (3) (55%) and the 8-substituted by-product comprising 8-nitro-5-methyl-4H-benzo[1,4]oxazin-3-one (4) (22%). These isomers are separated with difficulty at this point and were carried
- 25 on to the next step as a mixture. To a mixture of (3) (1.93 g, 9.27mmol) and (4) (0.48g, 2.32 mmol) dissolved in a solution of MeOH (300mL) and THF

(300mL) under argon was added 10% palladium on carbon (1.20g). The resulting solution was subjected to H₂ at one atmosphere pressure. After 16h the catalyst was filtered off and the resulting solution was concentrated in vacuo and purified by flash chromatography on silica with 50% hexane and ethyl acetate to give 6-amino-5-methyl-4H-benzo[1,4]oxazin-3-one (5) (0.96 g) in 46% yield and 8-amino-5-methyl-4H-benzo[1,4]oxazin-3-one (6) (0.17 g) in 8% yield. (5) (1.20g, 6.74 mmol), imidazoline-2-sulfonic acid (2.02g, 13.48 mmol) and triethylamine (2.35mL, 16.85 mmol) were heated at reflux in anhydrous acetonitrile (50mL) under argon for 48h. At that time an additional amount of imidazoline-2-sulfonic acid (1.01g, 6.74 mmol) and triethylamine (1.41mL, 10.12 mmol) were added and the resulting mixture was stirred an additional 24h. This solution was concentrated in vacuo and the residues were taken up in a solution of CHCl₃/isopropyl alcohol (3:1) and washed successively with NaOH (1N) and brine, dried over MgSO₄ and concentrated in vacuo. The resulting foam was purified by flash chromatography on silica with 20% methanol saturated with ammonia in chloroform to give 6-(imidazolidin-2-ylideneamino)-5-methyl-4H-benzo[1,4]oxazin-3-one (7) (0.42g) as a foam in 27% yield along with 55% recovered starting material. The HCl salt was recrystallized from a mixture of ethanol and diethyl ether (EtOH/Et₂O) to give fine white needles.

¹H NMR (DMSO): 2.10 (s, 3H); 3.59 (s, 4H); 4.53 (s, 2H); 6.83 (d, J=8.6 Hz, 1H); 6.90 (d, J= 8.6 Hz, 1H); 8.07 (brs, 2H); 10.15 (vbrs, 1H); 10.42 (s, 1H).


(6) (0.222 g, 1.35mmol), imidazoline-2-sulfonic acid (0.223 g, 1.49mmol) and triethylamine (0.415 mL, 2.98mmol) were heated at 95°C in anhydrous acetonitrile (10 mL) in a sealed tube for 2h. At that time an additional

amount of imidazoline-2-sulfonic acid (0.112 g, 0.75mmol) was added and the reaction was continued for an additional 16 h. This solution was concentrated in vacuo and the residues were taken up in a solution of CHCl_3 /isopropyl alcohol (3:1) and washed successively with NaOH (2N) and brine, dried (MgSO_4) and concentrated in vacuo. The resulting oil was recrystallized from CHCl_3 to give pure 6-(imidazolidin-2-ylideneamino)-5-methyl-4H-benzo[1,4]oxazin-3-one (8) (F) (0.048 g) as a white powder in 15% yield along with 35% recovered starting material. To a slurry of (8), (0.08 g, 0.321mmol) in anhydrous THF (50 mL) under argon in a 3-neck round-bottom flask equipped with reflux condenser was added borane-dimethyl sulfide complex (0.48 mL, 0.936mmol). The mixture was heated at reflux until starting material was no longer observed via thin layer chromatography (3 h). The reaction mixture was cooled to room temperature and carefully quenched by the dropwise addition of methanol. The crude reaction mixture was concentrated in vacuo and purified by flash chromatography on silica using 20% methanol saturated with ammonia/chloroform to give imidazolidin-2-ylidene-(5-methyl-3,4-dihydro-2H-benzo[1,4]oxazin-8-yl)amine (9) (G) (0.03 g) as the HCl salt in 37% yield. ^1H NMR (CDCl_3): 2.07 (s, 3H); 3.46 (t, $J=4.3\text{Hz}$, 2H); 3.55 (s, 4H); 4.24 (t, $J=4.3\text{Hz}$, 2H); 5.60 to 5.95 (vbrs, 2H); 6.44 (d, $J=8.0\text{ Hz}$, 1H); 6.57 (d, $J=8.0\text{ Hz}$, 1H).

Example H

Procedure for Preparation of 4(5)-phenylsulfanyl-1H-imidazole :

34



Procedure -

1-(N,N-dimethylsulfamoyl)imidazole (1.5g, 8.6 mmol) was taken up in 28mL of THF. The solution was cooled to -78°C and n-BuLi (5.4mL, 8.6 mmol) added dropwise via syringe. After stirring at -78°C for 1h TBSCl (1.3g, 8.56 mmol) in 10mL of THF was added. The bath was removed and the reaction allowed to warm-up to rt. The reaction mixture was stirred overnight. The reaction mixture was cooled to -20°C and n-BuLi (5.4 mL, 8.6 mmol) added. After 45 min phenyldisulfide (1.9g, 8.6 mmol) in 8mL of THF was added. The reaction mixture was stirred at rt for 48h. The reaction mixture was quenched with saturated ammonium chloride and extracted with ethyl acetate. The organic layer was collected and washed with water and then brine. The solution was dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (2.5% EtOAc/hexane) afforded 2.8g (7.0 mmol) of 2-(t-butyldimethylsilyl)-5-phenylsulfanylimidazole-1-sulfonic acid dimethylamide (1) as a yellow color oil. The compound (1) (2.8g, 7.0 mmol) was dissolved in THF and the solution cooled to 0°C. TBAF (7.0mL, 7.0 mmol) was added dropwise to the solution. The reaction mixture was stirred overnight at rt. The next day the reaction was quenched with water and extracted with ethyl acetate.


- The organic layer was washed with water followed by brine. The solution was dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (50% EtOAc/hexane) afforded 474mg of 5-phenylsulfanylimidazole-1-sulfonic acid dimethylamide (2) and 290mg of
- 5 5-phenylsulfanyl-1H-imidazole (3) (H). The 478mg of (2) was added to 2N HCl and the solution heated at reflux for 2h. The reaction mixture was made basic with 2N NaOH and extracted with ethyl acetate. The organic layer was washed with water followed by brine. The solution was dried over sodium sulfate and the solvent removed under reduced pressure.
- 10 Flash chromatography (EtOAc) afforded (3) as a white crystalline solid. A combined total of 360mg (2.0 mmol) of (3) is recovered.
- ¹H NMR (300 MHz, CD₃OD) 7.91 (s, 1H), 7.37 (s, 1H), 7.19-7.23 (m, 2H), 7.07-7.11 (m, 3H).

15

Example I

Procedure for Preparation of 4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-4,5-dihydro-1H-imidazole, methane sulfonic acid salt :

36



Procedure -

To 1,2,3,4-tetrahydronaphthalene-2-carboxylic acid (1) (4.93g, 27.42 mmol) in anhydrous THF (250mL) at 20°C under argon was added 3.26 mL (32.90 mmol) borane-dimethylsulfide ($\text{BH}_3\text{-Me}_2\text{S}$) via syringe. After stirring for 16h MeOH (4mL) was added and the mixture was warmed to 55°C until no more gas was evolved. The mixture was concentrated to an

oil, taken up in Et₂O and washed successively with 2M phosphoric acid, saturated sodium bicarbonate, water and brine and then dried over MgSO₄ and reconcentrated. The resulting oil was purified by high vacuum Kugelrohr at 150°C to give pure alcohol (1,2,3,4-tetrahydronaphthalen-2-yl)methanol (2) (4.09g) in 93% yield. To triphenylphosphine (10.179g, 38.809 mmol) and imidazole (2.64g, 38.809 mmol) in anhydrous benzene (175mL) was added the iodine (8.60g, 33.865 mmol) in benzene (75mL) with rapid stirring followed by (2) in benzene (50mL). After 3h the solids were filtered off and the filtrate was reduced in vacuo to a volume of 50mL to which was added hexane (200mL). The resultant solids were filtered off and the filtrate was washed successively with water and brine, dried over MgSO₄ and concentrated in vacuo. The resulting oil was purified by flash chromatography on silica with hexane to give pure 2-iodomethyl-1,2,3,4-tetrahydronaphthalene (3) (6.239g) in 90% yield. To (3) (10.02 g, 36.85 mmol) and CuI (1.41g, 7.37 mmol) in anhydrous THF (50mL) at -78°C under argon was added vinylmagnesium bromide (1M in THF, 73.70mL, 73.70 mmol) slowly at a speed at which no color developed. This solution was allowed to warm to 0°C and stirred for 6h. The resulting mixture was recooled to -40°C and quenched by the careful addition of 2M phosphoric acid (35mL). This solution was diluted with 100mL water and extracted with hexanes. The organic fractions were washed successively with water and brine, dried over MgSO₄ and concentrated in vacuo. The resulting oil was purified by flash chromatography on silica with hexane to give 2-allyl-1,2,3,4-tetrahydronaphthalene (4) (5.618g) in 88% yield. (4) (5.615g, 32.645 mmol) and meta-chloroperbenzoic acid (m-CPBA) (14.08g, 81.613 mmol) were stirred in anhydrous methylene chloride (50mL) for 16h. The solids

were filtered off and potassium fluoride KF (5.11g, 88.142 mmol) was added and this mixture was stirred an additional hour. The solids were filtered off and the reaction was concentrated in vacuo. The resulting oil was purified by flash chromatography on silica with 5% ethyl acetate in hexane to give 2-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)oxirane (5) (5.41g) in 88% yield. To (5) (1.626g, 8.649 mmol) in a solution of acetone (20mL) and water (5mL) was added sodium azide (1.97g, 30.271 mmol). This solution was warmed to 85°C and stirred for 48h. The solution was concentrated in vacuo and the residues were taken up in CHCl_3 and washed successively with water and brine, dried over MgSO_4 and concentrated in vacuo. The resulting oil was purified by flash chromatography on silica with 30% ethyl acetate in hexane to give pure 1-azido-3-(1,2,3,4-tetrahydronaphthalen-2-yl)propan-2-ol (6) (1.762g) in 88% yield. A mixture of (6) (1.88g, 8.140 mmol), triphenylphosphine (2.67g, 10.173 mmol), phthalimide (1.50g, 10.173 mmol), diethyl azodicarboxylate (DEAD) (1.77g, 10.173 mmol) were stirred in anhydrous THF (50mL) for 4h. This solution was concentrated in vacuo, taken up in a solution of hexane (25mL) and ether (25mL) and stirred for 16h. The solids were filtered off and the filtrate was concentrated in vacuo. The resulting oil was purified by flash chromatography on silica with 20% ethyl acetate in hexane to give 2-[1-azidomethyl]-2-(1,2,3,4-tetrahydronaphthalen-2-yl)ethylisoindole-1,3-dione (7) (2.487g) contaminated with a small amount of impurity which was carried on without further purification. A mixture of (7) (3.93g, 10.917 mmol) and hydrazine (0.680mL, 21.833 mmol) were heated in ethanol (60mL) at reflux for 16h. The solids were filtered off and the filtrate was concentrated in vacuo. The residues were purified by flash

- chromatography on silica with 5% MeOH in CH_2Cl_2 to give 1-azidomethyl-2-(1,2,3,4-tetrahydronaphthalen-2-yl)ethylamine (8) (2.057g) in 88% yield. A mixture of (8) (2.056g, 8.940 mmol) and 10% palladium on carbon (0.260 g) were stirred in MeOH (30mL) under 1 atmosphere of hydrogen for 16h.
- 5 The solids were filtered off and the filtrate was concentrated in vacuo. The residues were purified by flash chromatography on silica with 10% ammonia saturated MeOH in CH_2Cl_2 to give 3-(1,2,3,4-tetrahydronaphthalen-2-yl)propane-1,2-dione (9) (1.557g) in 85% yield. A mixture of (9) (0.590g, 2.892 mmol) and methanesulfonic acid (0.980mL, 14.460 mmol)
- 10 were heated in triethylorthoformate (10mL) at 105°C 3h. The reaction was concentrated in vacuo and the solids were filtered off. Subsequent recrystallization of these solids from a mixture of MeOH and ether gave pure 4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-4,5-dihydro-1H-imidazole, methane sulfonic acid salt (I) (0.435g) in 48% yield.
- 15 ^1H NMR (CDCl_3): 1.37 to 1.56 (m, 1H); 1.56 to 1.70 (m, 1H); 1.80 to 2.02 (m, 2H); 2.32 to 2.55 (m, 2H); 2.72 (s, 3H); 2.75 to 2.95 (m, 3H); 3.48 to 3.59 (m, 1H); 3.93 to 4.08 (m, 1H); 4.31 to 4.47 (m, 1H); 7.00 to 7.20 (m, 4H); 8.46 (s, 1H); 10.04 (s, 1H); 10.35 (brs, 1H).


20

Example J-1

Procedure for Preparation of 4(5)-cyclohexylmethyl-1H-imidazole :

25

40



Procedure -

- 2-Tert-butyldimethylsilyl-1-dimethylsulfamoyl imidazole (1) (4.1g, 14.2 mmol) is taken up in 47 mL of anhydrous THF and cooled to -20°C . $n\text{-BuLi}$ (8.9 mL, 14.2 mmol) is added dropwise to the solution of (1). The resultant solution is stirred at -20°C for 45 min. Cyclohexylmethyl iodide (2) (3.14g, 14 mmol) is then added dropwise to the reaction mixture. Then reaction is warmed to rt and stirred overnight. The next day the reaction is quenched with saturated ammonium chloride and diluted with water. The mixture is extracted with ethyl acetate (3 x 100 mL). The organic layers are combined and washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (4:1 ethyl acetate/ hexane) affords 2.26g (5.6 mmol) of 5-cyclohexylmethyl-2-tert-butyldimethylsilyl-1-dimethylsulfamoyl imidazole (3). (3) (2.26g, 5.6 mmol) is taken up in 56 mL of THF and cooled to 0°C . A 1M solution of TBAF in THF (5.6 mL, 5.6 mmol) is added dropwise to the solution of (3). The reaction is warmed to rt and stirred overnight. The next day the reaction is quenched with water and then extracted with ethyl acetate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate

and the solvent removed under reduced pressure. Flash chromatography (1:1 ethyl acetate/ hexane) affords 1.2g (4.42 mmol) of 5-cyclohexylmethyl-1-dimethylsulfamoyl imidazole (4). (4) (1.2g, 4.42 mmol) is taken up in 25 mL of a 1.5N HCl solution and heated at reflux for 2h. The reaction is cool
5 to rt and diluted with ethyl acetate. The mixture is brought to pH 13 with 2N NaOH and then extracted with chloroform (4 x 100 mL). The organic layers are combined and washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (9:1 chloroform/ methanol)
10 affords 700 mg (4.27 mmol) of 4(5)-cyclohexylmethyl-1H-imidazole (5) (J-1).

¹H NMR (CDCl₃): 0.92 to 1.0 (m, 2H); 1.16 to 1.26 (m, 3H); 1.57 to 1.73 (m, 6H); 2.48 (d, J=6.9 Hz, 2H); 6.77 (s, 1H); 7.56 (s, 1H)

Example J-2

15 (S)-2-iodomethyl-1,2,3,4-tetrahydronaphthalene is substituted into the method of Example J-1 to yield (S)-4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole. (S)-2-iodomethyl-1,2,3,4-tetrahydronaphthalene was prepared from (S)-1,2,3,4-tetrahydro-2-naphthoic acid. (S)-1,2,3,4-tetrahydro-2-naphthoic acid was prepared from the resolution of 1,2,3,4-
20 tetrahydro-2-naphthoic acid (*J. Med. Chem.* 1983, 26, 328-334)


Example J-3

(R)-2-iodomethyl-1,2,3,4-tetrahydronaphthalene is substituted into the method of Example J-1 to yield (R)-4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole. (R)-2-iodomethyl-1,2,3,4-tetrahydronaphthalene
25 was prepared from (R)-1,2,3,4-tetrahydro-2-naphthoic acid. (R)-1,2,3,4-

tetrahydro-2-naphthoic acid was prepared from the resolution of 1,2,3,4-tetrahydro-2-naphthoic acid (*J. Med. Chem.* 1983, 26, 328-334)

Example K-1

- 5 Procedure for Preparation of 4(5)-(4,5,6,7-tetrahydrobenzo[b]thiophen-2-ylmethyl)-1H-imidazole :



Procedure -

- 4,5,6,7-tetrahydrobenzo[b]thiophene (1) (2.1g, 15 mmol) is taken up in 75mL of anhydrous THF and cooled to -78°C . $n\text{-BuLi}$ (6.0mL, 15 mmol) is added dropwise to the solution of (1). The resultant solution is stirred at -78°C for 60 min. 1-Dimethylsulfonyl-2-t-butyl-5-(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)-1H-imidazole (3) is then added to the solution.
- 15

imidazolecarboxaldehyde (2) (4.8g, 15 mmol) in 25mL of THF is added to the reaction. The reaction is warmed to rt and stirred for 2h before being quenched with water and diluted with ethyl acetate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (1:3 ethyl acetate/ hexane) affords 5.2g (11 mmol) of 2-(tert-butyl dimethylsilyl)-5-[hydroxy-(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]imidazole-1-sulfonic acid dimethylamide (3). (3) (5.2g, 11.3 mmol) is taken up in 57mL of THF. A 1M solution of tetra-n-butylammonium fluoride (TBAF) in THF (11.3mL, 11.3 mmol) is added dropwise to the solution of (3). The reaction is stirred for 1h 15min reaction before being quenched with water and then extracted with ethyl acetate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Recrystallization from hexane/ethyl acetate affords 5-[hydroxy-(4,5,6,7-tetrahydrobenzo[b]thiophen-2-yl)methyl]imidazole-1-sulfonic acid dimethylamide (4) (2.1g, 6.2 mmol). An additional 2g of the crude product is also recovered. (4) (2.0g, 5.9 mmol) is taken up in 78mL of dichloromethane, to the solution is added 7.5 mL (46.9 mmol) of triethylsilane and 14.4 mL (0.19 mol) of trifluoroacetic acid. The reaction is stirred at rt overnight and then quenched with water and neutralized with 2N NaOH. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography using a 1:1 mixture of ethyl acetate and hexane affords 0.75g (2.3 mmol) of 5-(4,5,6,7-tetrahydrobenzo[b]thiophen-2-ylmethyl)imidazole-1-sulfonic acid

dimethylamide (5). (5) (0.42g, 1.55 mmol) is taken up in 15mL of a 1.5N HCl solution and heated at reflux for 2h and then stirred at rt overnight. The reaction is diluted with ethyl acetate, neutralized with 2N NaOH. The organic layer is washed with water followed by brine. The organic phase
5 is dried over sodium sulfate and the solvent removed under reduced pressure. The crude product is dissolved in methanol and an excess of HCl in ether is added. Solvent is removed under reduced pressure to afford 0.6g (2.3 mmol) of 4(5)-(4,5,6,7-tetrahydrobenzo[b]thiophen-2-ylmethyl)-1H-imidazole (6) (K-1).

10 ¹H NMR (CD₃OD): 8.80 (s, 1H); 7.34 (s, 1H); 6.57 (s, 1H); 4.18 (s, 2H); 2.65 to 2.69 (m, 2H); 2.51 to 2.55 (m, 2H); 1.74 to 1.83 (m, 4H)

Example K-2

2-(Tert-butyl) furan is substituted into the method of Example K-1 to yield 4(5)-(5-tert-butylfuran-2-ylmethyl)-1H-imidazole

15


Example K-3

5,6-Dihydro-4H-thieno[2,3-b]thiopyran is substituted into the method of Example K-1 to yield 4(5)-(5,6-dihydro-4H-thieno[2,3-b]thiopyran-2-ylmethyl)-1H-imidazole

Example L

20 Procedure for Preparation of 4(5)-(1-furan-2-ylethyl)-1H-imidazole:

45



Procedure -

- 2-(Tert-butyldimethylsilyl)-1-(dimethylsulfamoyl)imidazole (**1**) (3.3 g, 11.4 mmol) is taken up in 38mL of anhydrous THF and cooled to -78°C . $n\text{-BuLi}$ (7.2mL, 11.4 mmol) is added dropwise to the solution of (**1**). The resultant solution is stirred at -78°C for 30 min. 2-Furfural (**2**) (0.94mL, 11.4 mmol) is added to the reaction. The reaction is warmed to rt and stirred

overnight. The next day the reaction is quenched with saturated ammonium chloride and diluted with ethyl acetate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (4:1 ethyl acetate/ hexane) affords 4.4g (11.4 mmol) of 2-(t-butyl dimethylsilyl)-5-(furan-2-ylhydroxy-methyl)imidazole-1-sulfonic acid dimethylamide (3). (3) (4.4g, 11.4 mmol) is taken up in 110mL of THF and cool to 0°C. A 1M solution of tetra-n-butylammonium fluoride (TBAF) in THF (11.4mL, 11.4 mmol) is added dropwise to the solution of (3). The reaction is stirred overnight at rt. The next day the reaction is quenched with water and then extracted with ethyl acetate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. 3.9g of crude 5-(furan-2-ylhydroxymethyl)imidazole-1-sulfonic acid dimethylamide (4) is recovered. (4) (1.0g, 3.7 mmol) is taken up in 37mL of dichloromethane, to the solution is added 1.6g (18.5 mmol) of manganese dioxide. The reaction is stirred at rt overnight and then filtered through celite. The eluent is collected and the solvent removed under reduced pressure. Flash chromatography using a 1:1 mixture of ethyl acetate and hexane affords 0.69g (2.6 mmol) of 5-(furan-2-ylcarbonyl)imidazole-1-sulfonic acid dimethylamide (5). (5) (0.69g, 2.6 mmol) is taken up in 26mL of THF. The solution is cool to -78°C. 1.7mL (5.1 mmol) of a 3M solution of methylmagnesium chloride is added. After stirring at -78° C for 1.5h reaction is warmed to rt and stirred for an additional hour. The reaction is quenched with water and then extracted with ethyl acetate. The organic layer is washed with water followed by

brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Crystallization from ether/hexane affords 0.39g (1.4 mmol) of 5-(1-furan-2-yl-1-hydroxyethyl)imidazole-1-sulfonic acid dimethylamide (6). An additional 0.19g of (6) is recovered.

5 (6) (0.58g, 2.0 mmol) is taken up in 27mL of dichloromethane, to the solution is added 2.6 mL (16.3 mmol) of triethylsilane and 5.5 mL (71.4 mmol) of trifluoroacetic acid. The reaction is stirred at rt overnight and then quenched with water and neutralized with solid sodium bicarbonate. The organic layer is washed with water followed by brine. The organic

10 phase is dried over sodium sulfate and the solvent removed under reduced pressure.

Flash chromatography using a 2:1 mixture of ethyl acetate and hexane affords 0.53g (2.0 mmol) of 5-(1-furan-2-ylethyl)imidazole-1-sulfonic acid dimethylamide (7). (7) (0.34g, 1.3 mmol) is taken up in 10mL of a 1.5N HCl


15 solution and heated at reflux for 30min and then stirred at rt overnight. The reaction is diluted with ethyl acetate and then made basic with 2N NaOH. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (10:1 chloroform/methanol)

20 affords 0.1g (0.62 mmol) of 4(5)-(1-furan-2-ylethyl)-1H-imidazole (8) (L).

¹H NMR (300 MHz, CDCl₃) 7.56 (m, 1H), 7.33-7.34 (m, 1H), 6.81 (m, 1H), 6.29-6.31 (m, 1H), 6.06-6.07 (m, 1H), 4.22 (q, J= 7.2 Hz, 1H), 1.63 (d, J= 7.2 Hz, 3H).

Example M

Procedure for Preparation of 4(5)-(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-4-methyl-1H-imidazole :



5

Procedure -

4-Methyl-1-(dimethylsulfamoyl)imidazole (1) (2.0g, 10.6 mmol) is taken up in 42mL of anhydrous THF and cooled to -78°C . $n\text{-BuLi}$ (6.6mL, 10.6 mmol) is added dropwise to the solution of (1). The resultant solution is stirred at -78°C for 30 min. Tert-butyldimethylsilylchloride (TBSCl) (1.6g, 10.6 mmol) in 10mL of THF is added to the reaction. The reaction is warmed to rt and stirred overnight. The next day the reaction is cooled to -20°C and 7.3mL (11.6 mmol) of $n\text{-BuLi}$ added. After stirring at -20°C for 30 min, 1,4-benzodioxan-6-carboxaldehyde (2) (1.92g, 11.7 mmol) in 10mL of

THF is added to the reaction mixture. Then reaction is warmed to rt and stirred for 3h. The reaction is quenched with water and diluted with ethyl acetate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography (1:2 ethyl acetate/ hexane) affords 3.9g (8.4 mmol) of 2-(t-butyl dimethylsilyl)-5-[(2,3-dihydro benzo[1,4]dioxin-6-yl)hydroxymethyl]-4-methylimidazole-1-sulfonic acid dimethylamide (3). (3) (1.0g, 2.14 mmol) is taken up in 21mL of THF. A 1M solution of tetra-n-butylammonium fluoride (TBAF) in THF (2.35mL, 2.35 mmol) is added dropwise to the solution of (3). The reaction is stirred for 30min at rt. The reaction is quenched with water and then extracted with ethyl acetate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography using ethyl acetate as eluant affords 0.75g (2.12 mmol) 5-[(2,3-dihydrobenzo[1,4]dioxin-6-yl)hydroxymethyl]-4-methylimidazole-1-sulfonic acid dimethylamide (4). (4) (0.75g, 2.12 mmol) is taken up in 28mL of dichloromethane, to the solution is added 2.7mL (17.0 mmol) of triethylsilane and 5.2mL (67.8 mmol) of trifluoroacetic acid. The reaction is stirred at rt overnight and then quenched with water and neutralized with solid sodium bicarbonate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure. Flash chromatography using a 3:1 mixture of ethyl acetate and hexane affords 0.63g (1.87 mmol) of 5-(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-4-methylimidazole-1-sulfonic acid dimethylamide (5). (5) (0.63g, 1.87 mmol) is taken up in 10mL of a 1.5N

HCl solution and heated at reflux for. The reaction is diluted with ethyl acetate, neutralized with solid sodium bicarbonate. The organic layer is washed with water followed by brine. The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure.


- 5 Crystallization from ether/hexane affords 0.33g (1.43 mmol) of 4(5)-(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-4-methyl-1H-imidazole (6) (M).
¹H NMR (300 MHz, acetone-d⁶) 7.37 (s, 1H), 6.66-6.67 (m, 3H), 4.18 (s, 4H), 3.73 (s, 1H), 2.13 (s, 3H)

10

Example N

Procedure for Preparation of 2-(3H-imidazol-4(5)-ylmethyl)-3,4,5,6,7,8-hexahydro-2H-naphthalen-1-one (N-1), 4(5)-(2,3,4,4a,5,6,7,8-octahydronaphthalen-2-ylmethyl)-1H-imidazole (N-2) and 4(5)-(1,2,3,4,5,6,7,8-octahydronaphthalen-2-ylmethyl)-1H-imidazole (N-3):

15



Procedure :

1-Decalone (10.0g, 66 mmol) and 4(5)-imidazole carboxaldehyde (6.3g, 66 mmol) were added to 100 mL of ethanol. To the solution was

added NaOH (5.2g, 130 mmol) in 20 mL of water. The reaction was heated at reflux for 5 days. The reaction was cooled to rt and made basic with aqueous HCl. The solution was extracted with THF/ethyl acetate. The organic layers were combined and washed with brine. The organic phase
5 was dried over magnesium sulfate and the solvent removed under reduced pressure to afford the crude product. The crude product was heated at reflux in 40% H_2SO_4 for 1 day. The reaction was cooled to rt and made basic with saturated K_2CO_3 . The solution was extracted with THF/ethyl acetate. The organic layers were combined and washed with brine. The
10 organic phase was dried over magnesium sulfate and the solvent removed under reduced pressure. Purification by flash chromatography (15:1 CH_2Cl_2 /MeOH) afforded N-1 (4.9g, 32% yield).

1H NMR : 7.55 (s,1H), 6.77 (s, 1H), 3.08-3.14 (m, 2H), 1.52-2.46 (m, 13H).


The free base of the hydrochloride salt of N-1 (3.0g, 11 mmol) was
15 generated with NaOH and then added to diethylene glycol (100mL). To the solution was added hydrazine hydrate (3.2 mL, 100 mmol) and the reaction was left to stir overnight at rt. NaOH (3.1g, 77 mmol) was added and the solution heated at reflux for 5 days. The reaction was cooled to rt and diluted with water. The solution was extracted with THF/ethyl
20 acetate. The organic layers were combined and washed with brine. The organic phase was dried over magnesium sulfate and the solvent removed under reduced pressure. Purification by flash chromatography (8:1 CH_2Cl_2 /MeOH) afforded N-2 (0.64g, 27% yield).

1H NMR : 7.58 (s,1H), 6.76 (s, 1H), 5.24 (d, J= 4.3 Hz, 1H), 0.91-2.58 (m,
25 16H).

N-2 (1.0g, 4.6 mmol) was added to 10 mL of concentrated HCl. The solution was stirred at rt for 30 min and then neutralized with K_2CO_3 . The solution was extracted with THF/ethyl acetate. The organic layers were combined and washed with brine. The organic phase was dried over magnesium sulfate and the solvent removed under reduced pressure. Purification by flash chromatography (15:1 CH_2Cl_2 /MeOH) afforded N-3. 1H NMR : 7.54 (s,1H), 6.74 (s, 1H), 2.45-2.52 (m, 3H), 1.46-1.97 (m, 14H).

Example O

10 Procedure for Preparation of 4(5)-octahydro pentalen-2-ylmethyl)-1H-imidazole, hydrochloride:



Procedure-

- A. Following the synthesis of White and Whitesell, *Synthesis* pp. 602-3 (1975), ether (10 mL) was added to a flame-dried flask cooled to 0°C and then kept under an argon atmosphere. Then n-butyl lithium (35 mL of 2.5 M solution in hexane, 2.2 equiv.) was added and subsequently diisopropyl amine (14 mL, 2.5 equiv.) was added slowly and the mixture was allowed to stir for 30 min. at 0°C. To this generated solution of lithium diisopropyl amide was added cyclooctene oxide (5.0 g, 1.0 equiv.). The mixture was stirred at rt for one day and then heated to reflux under argon atmosphere for 2 days. The reaction was quenched by addition of NH_4Cl . The solution was extracted with THF/EtOAc. The organic extracts were combined, washed with brine, dried over magnesium sulfate and concentrated in vacuo to afford a yellow brown oil which was the 1-hydroxy-octahydropentalene. The compound was used without further purification in the next step.
- B. The alcohol thus obtained (5.0 g, 1 equiv.) was dissolved in dichloromethane (200 mL) and to this solution was added pyridinium chlorochromate (13 g, 1.5 equiv.) and the mixture was stirred at rt for one day. The solution was then filtered through a short column of SiO_2 using diethyl ether as eluent. The obtained solution was concentrated in vacuo to afford a pale green-yellow oil which was used without further purification in the next step.
- C. The octahydro-pentalen-1-one (5.0 g, 1.0 equiv.) of the above step was added to 4(5)-imidazolecarboxaldehyde (3.8 g, 1.0 equiv.) and 40% H_2SO_4 (20 ml) and the mixture was maintained at 90°C for 3 days. The reaction was then quenched by addition of ammonium hydroxide and


extracted with tetrahydrofuran/ethyl acetate. The organic extracts were combined, washed with brine, dried over magnesium sulfate. The resulting aqueous layer was neutralized with HCl/ NH₄Cl. The aqueous layer was re-extracted as above and the combined organic fractions were concentrated in vacuo to afford an orange solid.

D. This orange solid was dissolved in ethanol to which palladium on carbon (0.5 g) was added. The reaction flask was placed under 40 psi of hydrogen for one day. The reaction solution was filtered through celite with more ethanol used as eluent. The solution was concentrated in vacuo to afford a yellow brown oil. Purification by column chromatography using 17:1 chloroform/methanol afforded the ketone product in a somewhat impure state.

E. The ketone functionality was then removed by addition of the product of the step above (8.2 g, 1.0 equiv.) to diethylene glycol (80 mL) and hydrazine hydrate (13.0 g, 1.0 equiv.). This mixture was stirred overnight and then potassium hydroxide (11.0 g, 5.0 equiv.) was added and the solution was heated under reflux for one day. The reaction solution was cooled to rt and washed with water. The solution was extracted with THF/EtOAc and the combined fractions were washed with brine, dried over magnesium sulfate and concentrated in vacuo to afford a yellow oil. The monohydrochloride salt was made by dissolving this oil in anhydrous ethanol saturated with HCl and heating.

Example P

25 Procedure for the preparation of 7-(3H-imidazol-4(5)-ylmethyl)-6,7-dihydro-5H-isquinolin-8-one (P-1) and 7-(3H-imidazol-4(5)-ylmethyl)-5, 6, 7, 8-tetrahydroisoquinoline (P-2)



Procedure:

- 25 A. 3,4-lutidine (21.4g, 1 equiv.) was dissolved in 200 mL of water at 20°C and potassium permanganate was added in 6.32g portions twice daily for 5 days (total 63.2g, 2 equiv.). After 5 days the solution was stored in the freezer, then thawed and filtered through celite. The resulting colorless solution was concentrated at

90°C on a rotary evaporator until a white solid was obtained. This solid was recrystallized from 5N HCl to give 9.56g of white crystals. NMR indicated a mixture of two regioisomers with the desired isomer being the major product.

5 B. These crystals were heated in anhydrous ethanol saturated with HCl gas under argon and at reflux for 6 h. Then ethanol was removed from the solution by rotary evaporation and the residue was taken up in 100 mL of water and the pH was adjusted to between 7 and 8 with solid sodium bicarbonate. The aqueous phase was extracted with diethyl ether (3X) and the combined organic fractions were washed with brine, dried over magnesium sulfate and then filtered and concentrated to give
10 a colorless oil (3.56g, 10.8% yield).

C. Diisopropylamine 2.84g, 1.3 equiv.) was added to n-BuLi (11.21 mL, 1.3 equiv.) in 100 mL of anhydrous THF under argon at -78°C via syringe to produce lithium diisopropylamide in situ. To this solution was added the product of B above (3.56g, 1 equiv.) in 20 mL of tetrahydrofuran, via syringe and the mixture
15 was stirred at -78°C for 20 min. At this point methyl acrylate (4.85 mL 2.5 equiv.) in 20 mL of tetrahydrofuran was added dropwise through a cannula. The solution was stirred another 2 h before quenching by addition of 40 mL of 10% potassium acetate. The solution was allowed to warm to 20°C and then was concentrated on a rotary evaporator. The aqueous residue was extracted three times with chloroform.
20 The combined fractions were washed with brine and dried over magnesium sulfate, filtered and concentrated to a black solid, which was stored under high vacuum. Chromatography on silica gel with hexanes / ethyl acetate (7/3 → 6/4) afforded 2.41g (58.2%) of the desired product which was used without further purification in the next step.

25 D. The material from Step C (0.48g, 1 equiv.) was dissolved in 1 mL of 6M HCl and heated at 105°C for 16 h after which time the solution was concentrated to a solid by rotary evaporation at 80°C. The residue was taken up in 2 mL of water and neutralized with solid sodium bicarbonate. The neutralized solution was extracted


with chloroform (3X) and the combined fractions were washed with brine, dried over magnesium sulfate and concentrated to a colorless oil. (0.456g 93.4%).

- E. The isoquinolone (1.91 g, 1 equiv.) obtained in step D above was heated with 4(5)-imidazolecarboxaldehyde 1.25g, 1. equiv.) at 110°C in 15 mL of 40% sulfuric acid for 30 h. The reaction mixture was stored for several days at 0° C under argon. The solution was then diluted with 20 mL of water and basified to pH 8.9 with NH₄OH. Solids were collected by filtration and dried with high vacuum. The product was a yellow solid (2.81g, 96.1%) comprising a mixture of both positional isomers at the exo double bond.
- 10 F. The product of E, above, was dissolved in 150 mL of methanol and to this solution Pd/C (.412g, 0.15 wt. equiv.) was added. The methanolic solution was then saturated with H₂ by repeated evacuations and H₂ back-fill iterations. The solution was stirred under 1 atm. pressure of H₂ for 20 h until TLC revealed that no unsaturated starting material remained. The solution was filtered through celite and
- 15 concentrated to an oil. Chromatography on silica using dichloromethane and methanol (9/1) recovered pure product (1.853g 6504 %) as a white foam. This was taken up in methanol to which fumaric acid (0.4817g, 1.5 equiv.) was added with warming to dissolve the solids. The solution was cooled slowly and off-white crystals (0.826g, 74%) were obtained, which are represented as the compound P-1.
- 20 P-2 was obtained by hydrazine reduction in the same manner as described in Step E of Example O above.

Example Q

- Procedure for the preparation of (Z)-6-(3H-imidazol-4(5)-ylmethylene)-7,8-dihydro-6H-quinolin-5-one (Q-1), (E)-6-(3H-imidazol-4(5)-ylmethylene)-7,8-dihydro-6H-quinolin-5-one (Q-2), 6-(3H-imidazol-4(5)-ylmethyl)-7,8-dihydro-6H-quinolin-5-one (Q-3), 6-(3H-imidazol-4(5)-ylmethyl)-5,6,7,8-tetrahydroquinoline, dihydrochloride (Q-4) and 6-(3H-imidazol-4(5)-ylmethyl)-octahydroquinolin-5-one (Q-5)
- 25

58



progressing to 95/5 dichloromethane:methanol as eluent. The amidophosphonate intermediate was obtained in 2.139 g, 65.1%, yield.

C. The amidophosphonate was dissolved in 100 mL of anhydrous o-xylene and then 10% Pd / C was added with stirring. Freshly distilled acrolein was then added to the mixture via syringe and heated to reflux for 4 h, after which time the remaining acrolein was added and heating under reflux was continued for 44 h under a finger condenser and under argon. At that time TLC indicated some intermediate remained, so 0.5g addition Pd/ C was added and the mixture again was heated to reflux for another 8 h. The mixture was cooled to rt, filtered and concentrated on a rotovap to eliminate excess acrolein, until about 100 mL of o-xylene solution remained. This solution was cooled by addition of ice, and was extracted three times with 1N HCl. The combined aqueous fractions were extracted 3X with Et₂O. The aqueous phase was then cooled to 0°C and the pH was adjusted to ~10 using concentrated NaOH. The aqueous was then extracted 5X with 100 mL portions of chloroform. The combined chloroform fractions were washed with water and then brined and dried over magnesium sulfate, filtered, and finally concentrated to give 3.51 g of an oil in 84.4% yield of 7,8-dihydro-6H-quinolin-5-one.

D. The 4(5)-imidazole carboxaldehyde was condensed with the quinolinone as described in Step E of Example P and was obtained both Q-1 and Q-2.

E. The exo double bond was then reduced with palladium on carbon as described in Step F of Example P above to yield two products which were separated by chromatography to give Q-3 and A.

F. The keto group was removed by the same hydrazine reduction procedure as that described in Step E of Example O above to give Q4.


G. The fully-reduced quinoline ring product Q-5 was obtained by a standard reduction of A with lithium/ammonia. (Li, 10 equiv., in NH₃ at -78°C for 10 min, quenched with NH₄OH, gradual warming with NH₃ evaporation).

Procedure:

- A. The reactive azido reagent of the first step was generated in situ by addition of iodine monochloride (67.6 g, 1.15 equiv.) in 50 mL of acetonitrile dropwise through a dropping funnel to a stirred slurry of sodium azide (58.84 g, 2.5 equiv.) in 350 mL of anhydrous acetonitrile at -10°C and under argon. Addition was complete in 30 min, the mixture was stirred an additional 30 min and cyclohexenone (34.81 g, 1.0 equiv.) was added via a syringe and then stirred at 20°C for an additional 20 h. The mixture was then poured into a liter of water and extracted with three 200 mL portions of diethyl ether. The combined fractions were washed with 5% sodium thiosulfate solution and then brine. The organic phase was dried over magnesium sulfate, filtered and concentrated in vacuo at 20°C . The residues were taken up in 1 L of DMSO at 0°C and a second portion of NaN_3 was added and the mixture stirred while warming to ambient temperature. This mixture was then diluted with 2.5 L of ice water and extracted ten times with dichloromethane (10 X 250 mL). The combined organic fractions were concentrated on a rotovap to a volume of ~ 1 L and this concentrate was extracted three times with 250 mL of water, and then brine, and then dried over magnesium sulfate and concentrated to a dark oil (39.5 g) and stored at -40°C .
- The oil was purified by chromatography on silica using 9/1 to 8/2 hexane:ethyl acetate. Two isomers were recovered, the first with the azido group α to the ketone function was obtained in 13.22 g, 26.6%, yield. The β -isomer was obtained in 15.825 g, 32.0%, yield.
- B. Triphenyl phosphine was dissolved in 20 mL of dichloromethane and placed under an argon atmosphere at 20°C . The β -isomer obtained as described above was added via cannula to the stirred solution and maintained at 20°C for 2 h. As the reaction progressed nitrogen was liberated from the solution, and after 2 h TLC demonstrated there was no starting material remaining. The solution was concentrated and passed through a silica gel column with dichloromethane

Example R-1

Procedure for the preparation of (E)-6-(3H-imidazol-4(5)-ylmethylene)-7, 8-dihydro-6H-quinoxalin-5-one



Procedure:

A. A mixture of 5,6,7,8-tetrahydroquinoxaline (23.75g, 1 equiv.),
15 benzaldehyde (19.81 mL, 1.1 equiv.) and acetic anhydride (33.4 mL, 2.0 equiv.)
was stirred at 150°C under argon for 15 hr, after which time TLC indicated mostly
desired product with some starting materials remaining. Starting materials were
removed by vacuum distillation using a Vigreux column at 170°C. The pot residue
was then subjected to Kugelrohr distillation from 170 – 220°C. The first fraction
20 was slightly contaminated with starting materials (4.71g). A second fraction was
pure (18.93g). After applying high vacuum to the first fraction it crystallized.
Combined fractions yielded 20.11g, 51%.

B. The product from A, above, was dissolved in 100 mL of methanol and
warmed slightly, then cooled to –35 to –40°C and ozone was bubbled through the
25 solution. After a few minutes the starting material began to crystallize out of
solution and the solution was warmed and another 200 mL of methanol was added
and then the reaction was resumed. After about 30 minutes the solution turned pale
blue. Nitrogen was then introduced by bubbling through the solution for 30

minutes, then methyl sulfide (3.5 mL) was injected into the solution, whereafter the solution was stirred for another 30 min. at -35°C , then allowed to warm to ambient temperature with stirring. After about 48 hr. at 20°C the mixture was steam distilled to remove solvents to provide a residue of 8.4g of a yellow-brown oil.

- 5 This residue was taken up in diethyl ether and extracted 3x with 25 mL portions of 1N HCl. The combined aqueous fractions were washed with diethyl ether 3x. The aqueous solution was gradually basified to a pH of 8 with concentrated NaOH. The free amine was then extracted from the aqueous phase with chloroform (3x). The combined chloroform extracts were washed twice with brine, dried of MgSO_4 and
10 concentrated to a yellow oil (3.01g) After keeping under high vacuum for 1 hr., 2.97g remained. This was recrystallized from diethyl ether to give 2.35g of a bright yellow solid. Yield 67.5%.

- C. The 7,8-dihydroquinoxalin-5-one and 4(5)-imidazolecarboxaldehyde (Aldrich Chemicals) were suspended in 75 mL of anhydrous tetrahydrofuran at
15 20°C under argon followed by addition of piperidine followed by acetic acid. The mixture was stirred 16 h at 20°C . After 20 h, no traces of the quinoxalone remained as indicated by TLC. The solids were collected by filtration and washed with a small amount of tetrahydrofuran, followed by chloroform. The solid was dried under high vacuum to give 6.85g of R-1. Yield 90.3%.

20

Example R-2 and R-3

- In a similar manner as R-1, 5,6,7,8-tetrahydroisoquinoline (5.42g, 1 equiv., Aldrich) was stirred with benzaldehyde (5.182 g, 1.2 equiv.) and acetic anhydride (6.309 g, 2.0 g) which was vacuum distilled and used without further purification
25 in the next step. Yield (impure): 8.28 g.

The crude product (7.96 g) from the step above was subjected to ozonolysis as described in Step B above. After work-up and chromatography there was obtained 5.18 g of a pale oil. Yield: 97.8% assuming pure starting material.

The resulting 7,8-dihydro-6H-isoquinolin-5-one (1.692 g, 1 equiv.) was condensed with 4(5)-imidazolecarboxaldehyde as described in Step C above to yield 2.23 g of the unsaturated compound analogous to R-1 in the scheme above in 92.8% yield. This product was treated with palladium on carbon as described in Step F of

5 Example P to reduce the exo double bond to produce 6-(3H-imidazol-4(5)-ylmethyl)-7,8-dihydro-6H-isoquinolin-5-one (R-2) in 52%.

The ketone above was reduced using hydrazine and converted to the fumarate salt as detailed in Example P, Step F. Yield for the reduction: 62%. Yield of fumarate salt after recrystallization: 30.4% of 6-(3H-imidazol-4(5)-ylmethyl)-5,6,7,8-

10 tetrahydroisoquinoline (R-3).

Example S

A method for measuring α -agonist selectivity comprises the RSAT (Receptor Selection and Amplification Technology) assay as reported in

15 Messier et al. (1995) "High throughput assays of cloned adrenergic, muscarinic, neurokinin and neurotrophin receptors in living mammalian cells", *Pharmacol. Toxicol.* 76:308-11 and adapted for use with α_2 receptors. The assay measures a receptor-mediated loss of contact inhibition that results in selective proliferation of receptor-containing cells

20 in a mixed population of confluent cells. The increase in cell number is assessed with an appropriate transfected marker gene such as b-galactosidase, the activity of which can be easily measured in a 96-well format. Receptors that activate the G protein, G_q , elicit this response. α_2 receptors, which normally couple to G_i , activate the RSAT response

25 when coexpressed with a hybrid G_q protein that has a G_i receptor recognition domain, called $G_{q/i52}$. See Conklin et al. (1993) "Substitution

of three amino acids switches receptor specificity of $G_{\alpha a}$ to that of $G_{\alpha i}$."


Nature 363:274-6.





NIH-3T3 cells are plated at a density of 2×10^6 cells in 15 cm dishes and maintained in Dulbecco's modified Eagle's medium supplemented with 10% calf serum. One day later, cells are cotransfected by calcium phosphate precipitation with mammalian expression plasmids encoding p-SV-b-galactosidase (5-10 mg), receptor (1-2 mg) and G protein (1-2 mg). 40 mg salmon sperm DNA may also be included in the transfection mixture. Fresh media is added on the following day and 1-2 days later, cells are harvested and frozen in 50 assay aliquots. Cells are thawed and 100 μ l added to 100 μ l aliquots of various concentrations of drugs in triplicate in 96-well dishes. Incubations continue 72-96 hr at 37°. After washing with phosphate-buffered saline, b-galactosidase enzyme activity is determined by adding 200 μ l of the chromogenic substrate (consisting of 3.5 mM o-nitrophenyl-b-D-galactopyranoside and 0.5% nonidet P-40 in phosphate buffered saline), incubating overnight at 30° and measuring optical density at 420 nm. The absorbance is a measure of enzyme activity, which depends on cell number and reflects a receptor-mediated cell proliferation. The EC_{50} and maximal effect of each drug at each α_1 receptor is determined. The efficacy or intrinsic activity is calculated as a ratio of the maximal effect of the drug to the maximal effect of a standard full agonist for each receptor subtype. Brimonidine, also called UK14,304-18, is used as the standard agonist for the α_{2A} and α_{2C} receptors. Oxymetazoline is the standard agonist used for the α_{2B} receptor.





Table 1, below, provides the intrinsic activity values at subtypes of the α_2 -adrenoreceptor as determined in the RSAT assay for the compounds






of above Examples B through R and certain adrenergic compounds not having selective agonist activity at the $\alpha 2B$ or $\alpha 2B/\alpha 2C$ subtype(s). At the $\alpha 2A$ subtype, the compounds of the Examples are inactive or exhibit low efficacy (≤ 0.4). They have greater efficacy at the $\alpha 2B$ and the $\alpha 2C$ -subtypes than the $\alpha 2A$ -subtype. Therefore, unlike ophthalmic $\alpha 2$ -adrenoreceptor compounds such as clonidine and brimonidine, the compounds of Examples B through R can selectively activate $\alpha 2$ -adrenoreceptor subtypes other than the $\alpha 2A$ -subtype.






Table 1: Intrinsic Activity Relative to Brimonidine/Oxymetazoline





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
	oxymetazoline	0.63	1.0	0.58
	clonidine	0.78	0.75	0.55
	brimonidine	1.0	0.93	1.0
	4(5)-(3-methyl-thiophen-2-ylmethyl)-1H-imidazole	0.43	1.4	0.5
D-3	 bicyclo[2.2.1]hept-2-yl oxazolidin-2-ylidene amine	0	0.4	0





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>D-1</u>	 oxazolidin-2-ylidene-(3-phenylbicyclo[2.2.1]hept-2-yl) amine	0	0.47	0
<u>F</u>	 6-(imidazolidin-2-ylideneamino)-5-methyl-4H-benzo[1,4]oxazin-3-one	0.3	0.9	0.2
<u>G</u>	 imidazolidin-2-ylidene-(5-methyl-3,4-dihydro-2H-benzo[1,4]oxazin-8-yl) amine, hydrogen chloride salt	0.1	0.87	0.33
<u>J-1</u>	 4(5)-cyclohexylmethyl-1H-imidazole	0.1	0.83	0





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
	imidazole			
<u>E-1</u>	 <p>imidazolidin-2-ylidene-(4-methyl-3,4-dihydro-2H-benzo[1,4]oxazin-6-yl) amine</p>	0.33	0.83	0.35
<u>M</u>	 <p>4(5)-(2,3-dihydrobenzo[1,4]dioxin-6-ylmethyl)-4-methyl-1H-imidazole</p>	0.2	0.97	0.27
<u>C-2</u>	 <p>4(5)-thiophen-2-ylmethyl-1H-imidazole</p>	0.23	1.3	0.5
<u>C-1</u>	 <p>4(5)-thiophen-3-ylmethyl-1H-imidazole</p>	0	0.83	0





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>C-9</u>	 4(5)-benzo[b]thiophen-2-ylmethyl-1H-imidazole	0.06	0.88	0.43
<u>C-3</u>	 4(5)-(5-methylthiophen-2-ylmethyl)-1H-imidazole	0.1	0.88	0.43
<u>C-8</u>	 4(5)-benzyl-1H-imidazole	0.3	0.9	0.4
<u>H</u>	 4(5)-phenylsulfanyl-1H-imidazole	0.2	0.93	0.15
<u>C-5</u>	 4(5)-furan-2-ylmethyl-1H-imidazole	0	1.1	0.4





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>B-3b</u>	 <p>4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole</p>	0	0.7	0
<u>I-2</u>	 <p>(S)-4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole</p>	0	0.8	0
<u>I-3</u>	 <p>(R)-4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole</p>	0.1	1	0.15
<u>L</u>	 <p>4(5)-(1-furan-2-ylethyl)-1H-imidazole</p>	0.23	0.9	0.57
<u>C-6</u>	 <p>4(5)-furan-3-ylmethyl-1H-imidazole</p>	0.2	0.67	0.1





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>C-4</u>	 4(5)-(5-chlorothiophen-2-ylmethyl)-1H-imidazole	0.05	0.82	0.5
<u>D-2</u>	 oxazolidin-2-ylidene-(3-o-tolyl bicyclo[2.2.1]hept-2-yl) amine	0.25	0.75	0
<u>C-10</u>	 4(5)-benzofuran-2-ylmethyl-1H-imidazole	0.05	0.48	0.1
<u>C-7</u>	 4(5)-(5-methylfuran-2-ylmethyl)-1H-imidazole	0.08	0.73	0.2





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>B-3a</u>	 <p>2-(3H-imidazol-4(5)-ylmethyl)-3,4-dihydro-2H-naphthalen-1-one</p>	0.1	0.8	0.07
<u>I</u>	<p>CH₃SO₃H</p>  <p>4(5)-(1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-4,5-dihydro-1H-imidazole, methane sulfonic acid salt</p>	0	0.5	0.2
<u>B-2a</u>	 <p>3-(3H-imidazol-4(5)-ylmethylene)chroman-4-one</p>	0	0.63	0.15
<u>B-2b</u>	 <p>3-(3H-imidazol-4(5)-ylmethyl)chroman-4-one</p>	0	0.77	0






Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>B-2d</u>	 4(5)-chroman-3-ylmethyl-1H-imidazole	0	0.6	0
<u>B-2c</u>	 3-(3H-imidazol-4(5)-ylmethyl)chroman-4-ol	0	0.65	0
<u>B-9a</u>	 4(5)-(4,5,6,7-tetrahydrobenzo[b]thiophen-5-ylmethyl)-1H-imidazole	0.08	0.46	0
<u>B-4a</u>	 4(5)-(4-methyl-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole	0	0.75	0.1





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>B-4b</u>	 <p>2-(3H-imidazol-4(5)-ylmethyl)-4-methyl-3,4-dihydro-2H-naphthalen-1-one</p>	0.3	0.7	0.6
<u>B-11b</u>	 <p>6-(1H-imidazol-4(5)-ylmethylene)-6,7,8,9-tetrahydrobenzocyclohepten-5-one</p>	0	0.3	0
<u>B-6</u>	<p>HCl</p>  <p>4(5)-thiochrom-3-ylmethyl-1H-imidazole, hydrogen chloride salt</p>	0	0.35	0
<u>B-5b</u>	 <p>3-(3H-imidazol-4(5)-ylmethyl)thiochroman-4-one</p>	0	0.5	0.2




Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
B-5a	 <p>3-(3H-imidazol-4(5)-ylmethylene)thiochroman-4-one</p>	0	0.5	0.37
B-7a	 <p>2-(3H-imidazol-4(5)-ylmethylene)indan-1-one</p>	0	0.3	0
B-11a	 <p>4(5)-(6,7,8,9-tetrahydro-5H-benzocyclohepten-6-ylmethyl)-1H-imidazole</p>	0.4	0.9	0
B-7b	 <p>2-(3H-imidazol-4(5)-ylmethyl)indan-1-one</p>	0	0.3	0





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>B-1</u>	 <p>4(5)-(7-methoxy-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole, hydrogen chloride salt</p>	0.15	0.45	0.3
<u>B-1a</u>	 <p>2-(3H-imidazol-4(5)-ylmethyl)-7-methoxy-3,4-dihydro-2H-naphthalen-1-one</p>	0.15	0.6	0
<u>B-9b</u>	 <p>5-(3H-imidazol-4(5)-ylmethyl)-6,7-dihydro-5H-benzo[b]thiophen-4-one, hydrogen chloride salt</p>	0	0.68	0.15
<u>B-7c</u>	 <p>4(5)-indan-2-ylmethyl-1H-imidazole</p>	0	0.9	0





Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>B-10</u>	 <p>4(5)-(4,4-dimethyl-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole</p>	0	0.3	0
<u>B-8b</u>	 <p>4(5)-(7-methyl-1,2,3,4-tetrahydronaphthalen-2-ylmethyl)-1H-imidazole, hydrogen chloride salt</p>	0	0.6	0.2
<u>B-8a</u>	 <p>2-(3H-imidazol-4(5)-ylmethyl)-7-methyl-3,4-dihydro-2H-naphthalen-1-one</p>	0	0.4	0
<u>K-1</u>	 <p>4(5)-(4,5,6,7-tetrahydrobenzo[b]thiophen-2-ylmethyl)-1H-imidazole</p>	0	0.53	0


Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>C-12</u>	 4(5)-(4-bromothiophen-2-ylmethyl)-1H-imidazole	0.2	1.3	0.3
<u>C-13</u>	 4(5)-(4-phenylthiophen-2-ylmethyl)-1H-imidazole	0	0.5	0
<u>K-3</u>	 4(5)-(5,6-dihydro-4H-thieno[2,3-b]thiopyran-2-ylmethyl)-1H-imidazole	0	0.37	0
<u>K-2</u>	 4(5)-(5-tert-butylfuran-2-ylmethyl)-1H-imidazole	0	0.7	0
<u>C-11</u>	 4(5)-(5-ethylfuran-2-ylmethyl)-1H-imidazole	0.2	0.5	0

Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>C-14</u>	 <p>4(5)-(4-methylthiophen-2-ylmethyl)-1H-imidazole, hydrochloride salt</p>	0.27	0.7	0.3
<u>N-1</u>	 <p>2-(3H-imidazol-4(5)-ylmethyl)-3,4,5,6,7,8-hexahydro-2H-naphthalen-1-one, hydrochloride salt</p>	0.24	0.75	0.26
<u>Q-3</u>	 <p>6-(3H-imidazol-4(5)-ylmethyl)-7,8-dihydro-6H-quinolin-5-one</p>	0.1	0.9	0.23
<u>Q-2</u>		0.1	0.87	0.13

Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
	(E)-6-(3H-imidazol-4(5)-ylmethylene)-7,8-dihydro-6H-quinolin-5-one			
<u>Q-1</u>	 (Z)-6-(3H-imidazol-4(5)-ylmethylene)-7,8-dihydro-6H-quinolin-5-one	0	0.75	0.2
<u>N-2</u>	 4(5)-(2,3,4,4a,5,6,7,8-octahydronaphthalen-2-ylmethyl)-1H-imidazole	0	0.5	0.05
<u>Q-4</u>	 6-(3H-imidazol-4(5)-ylmethyl)-5,6,7,8-tetrahydroquinoline, dihydrochloride	0.1	0.8	0.1

Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>Q</u>	 <p>4(5)-octahydro pentalen-2-ylmethyl-1H-imidazole, hydrochloride</p>	0	0.67	0.1
<u>B-9c</u>	 <p>5-(octahydro benzo[b]thiophen-5-ylmethyl)-1H-imidazole, hydrochloride</p>	0	0.3	0
<u>R-3</u>	 <p>6-(3H-imidazol-4(5)-ylmethyl)-5,6,7,8-tetrahydro-isoquinoline, fumarate</p>	0	0.6	0.4
<u>R-2</u>	 <p>6-(3H-imidazol-4(5)-ylmethyl)-7,8-dihydro-6H-isoquinolin-5-one, dihydrochloride</p>	0	0.6	0.4

Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
<u>R-1</u>	 <p>(E)-6-(3H-imidazol-4(5)-ylmethylene)-7,8-dihydro-6H-quinoxalin-5-one</p>	0.3	0.8	0.4
<u>P-1</u>	 <p>7-(3H-imidazol-4(5)-ylmethyl)-6,7-dihydro-5H-isoquinolin-8-one, fumarate</p>	0	0.4	0
<u>P-2</u>	 <p>7-(3H-imidazol-4(5)-ylmethyl)-5,6,7,8-tetrahydroisoquinoline, fumarate</p>	0	0.4	0
<u>N-3</u>		0	0.75	0

Example	Structure/Compound	Brimonidine Alpha 2A	Oxymetazoline Alpha 2B	Brimonidine Alpha 2C
	4(5)-(1,2,3,4,5,6,7,8-octahydronaphthalen-2-ylmethyl)-1H-imidazole, fumarate			
<u>Q-5</u>	 6-(3H-imidazol-4(5)-ylmethyl)-octahydroquinolin-5-one	0	1.0	0

Example T

IOP-Lowering and Sedative Side Effects

5

Measurements of IOP were made in fully conscious female cynomolgus monkeys weighing 3-4 kg with sustained elevated IOP that was produced in the right eye by argon laser photocoagulation of the trabecular meshwork. Animals were usable for experiments ~ 2 months following surgery. During the experiments, monkeys sat in specially designed chairs (Primate Products, San Francisco), and were fed orange juice and fruit as needed. A 30R model Digilab pneumatonometer (Alcon, Texas) was used to measure IOP.

10

Twenty five µl of an anesthetic (proparacaine) was topically applied to each monkey before IOP measurements to minimize ocular discomfort due to tonometry. Two baseline measurements were made prior to instillation of the drugs, followed by periodic measurements up to 6 hours

15

post-instillation. The test compounds were administered unilaterally as a single 50 μ l eye drop; the contralateral eyes received an equal volume of saline.

Many of the α 2B or α 2B/2C selective compounds of the examples
5 were tested in the monkeys. Surprisingly, as Table 2 shows, these structurally diverse compounds all lowered IOP in the treated eye.

At the same time, sedation was measured and assessed according to the following score: 0 = alert, typical vocalization, movement, etc.; 1 = calm, less movement; 2 = slightly sedated, some vocalization, responsive to
10 stimulation; 3 = sedated, no vocalization, some response to stimulation; 4 = asleep.

The compounds of the present invention also did not cause sedation. This contrasts with the action of clonidine and brimonidine, which caused sedation.

15

Table 2. The effects of α 2-adrenoceptor agonists on IOP and sedation in conscious cynomolgus monkeys following ocular administration in eyes made unilaterally hypertensive by argon laser photocoagulation. Measurements were made periodically up to 6 hours. Sedation was assessed subjectively during the IOP
20 experiments using the following scoring: 0 = alert, typical vocalization, movement, etc.; 1 = calm, less movement; 2 = slightly sedated, some vocalization, responsive to stimulation; 3 = sedated, no vocalization, some response to stimulation; 4 = asleep. Number of animals per group = (6-9).

25

Table 2	Maximum % Decrease From Pretreatment Levels		
	Compounds	Dose (%)	Hypertensive Eye Sedation (0-4)
	Saline	-	7 ± 2 0-1
	Clonidine	0.1	25 ± 4 1
		0.3	41 ± 5 2
	Brimonidine	0.1	25 ± 3 1
		0.3	40 ± 4 2
	J-1	1	26 ± 5 0
		3	33 ± 3 0
	E-1	0.3	25 ± 4 0
		1	27 ± 3 0
	C-1	1	25 ± 4 0
		3	29 ± 4 0
	D-1	1	25.6 ± 3.9 0
	M	1	22.5 ± 5.4 0
	C-2	1	29.6 ± 5.5 0

C-9	0.3	13.7 ± 4.5	0
	1	25.1 ± 4.9	0
C-3	0.3	20.6 ± 4.8	0
	1	25.0 ± 6.4	0
C-8	1	31.2 ± 3.3	0
B-3b	0.1	25.9 ± 3.5	0
	0.3	31.2 ± 4.3	0
C-4	0.3	17.7 ± 4.0	0
	1	29.3 ± 4.9	0
C-7	1	32.3 ± 5.7	0
J-2	0.03	12.4 ± 3.7	0
	0.3	27.3 ± 3.1	0
J-3	0.03	16.4 ± 4.7	0
	0.3	26.5 ± 3.8	0
B-2d	0.1	22.0 ± 4.6	0
	0.3	17.0 ± 4.2	0
	1	18.1 ± 5.2	0

B-9a	0.03	17.6 ± 1.7	0
	0.1	26.7 ± 6.1	0
	0.3	24.8 ± 3.3	0
	1	26.8 ± 5.4	0
B-6	0.3	13.8 ± 2.4	0
	1	22.1 ± 6.3	0
B-9b	0.1	18.7 ± 5.5	0
	0.3	26.9 ± 6.1	0

Example U

Measurement of Cardiovascular Side Effects

5 Cardiovascular measurements were made in a different group of monkeys using a BP 100S automated sphygmomanometer (Nippon Colin, Japan). Intravenous (IV) administration of certain of the compounds of the present invention at doses ten to thirty times higher than the doses for clonidine and brimonidine did not reduce heart rate or lower blood
 10 pressure. Interestingly, the compound 4(5)-3-methylthiophen-2-ylmethyl)-1H-imidazole, which has intrinsic activity of 0.43 at the α_2A -subtype, exhibited a weak effect on heart rate. Clonidine and brimonidine had even greater effects on heart rate. See Table 3 below.

15 **Table 3.** The effects of α_2 -adrenoceptor agonists on cardiovascular variables in conscious cynomolgus monkeys following i.v. administration. Measurements were made periodically up to 6 hours. Number of animals per group = (6-10).

Table 3		Maximum % Decrease From Pretreatment Levels	
Compounds	Dose ($\mu\text{g/kg}$)	Mean Arterial Blood Pressure	Heart Rate
Saline	-	7 ± 4	8 ± 3
Clonidine	17	29 ± 7	32 ± 4
	50	35 ± 5	50 ± 5
Brimonidine	17	36 ± 3	52 ± 3
	50	37 ± 5	54 ± 3
J-1	17	7 ± 5.3	13 ± 4
	50	4 ± 2	6 ± 2
	167	7 ± 5	3 ± 3
	500	13 ± 3	7 ± 4
E-1	17	7 ± 4	11 ± 4
	50	7 ± 2	14 ± 5
	167	9 ± 4	11 ± 5
C-1	50	12.8 ± 12	12 ± 4
	500	$+5 \pm 8^*$	$+11 \pm 9^*$
M	500	0.8 ± 2.3	5.5 ± 1.9
C-2	500	6.6 ± 1.7	6.5 ± 2.9
C-9	3.0	5.0 ± 2.3	9.4 ± 3.0
	17	1.0 ± 4.1	$+9.4 \pm 1.8^*$
	50	0.1 ± 3.8	16 ± 3.2
	500	6.0 ± 2.2	5.9 ± 3.3
C-3	500	2.3 ± 2.7	10.6 ± 3.4
C-8	500	5.5 ± 2.7	16.6 ± 1.9
C-5	500	3.9 ± 2.8	7.1 ± 3.9
B-3b	50	2.4 ± 4.3	10.0 ± 2.8
C-4	500	5.3 ± 2.9	10.9 ± 3.6

C-7	500	3.0 ± 3.9	6.1 ± 3.7
J-2	500	$+0.6 \pm 3.1^*$	6.4 ± 3.3
J-3	500	$+1.0 \pm 2.1^*$	$+10.6 \pm 6.0^*$
B-2b	500	5.7 ± 1.4	6.4 ± 3.6
B-2d	500	$+8.9 \pm 3.4^*$	$+15.5 \pm 3.4^*$
B-9a	500	$+10.8 \pm 3.2^*$	$+23.8 \pm 4.4^*$
B-9b	500	2.8 ± 1.8	$+20.2 \pm 3.4^*$
4(5)-(3-methylthiophen-2-ylmethyl)-1H-imidazole	50	9 ± 3	23 ± 4
	167	8 ± 6	32 ± 8
* showed increase from base levels			

EXAMPLE V

The studies in the above Examples T and U demonstrate that a therapeutic effect of alpha2 agonists can be separated from sedative and cardiovascular side effects. This separation is accomplished with compounds that share the property of being preferentially active at the alpha2B and alpha2B/alpha2C subtypes relative to the alpha2A subtype.

The prior art alpha2 adrenergic agonists, which activate all three alpha2 receptors, cause sedation, hypotension and bradycardia, preventing or severely limiting their use for treating diseases and disorders that are known to be ameliorated by them. Such diseases and disorders include muscle spasticity including hyperactive micturition, diarrhea, diuresis, withdrawal syndromes, pain including neuropathic pain, neurodegenerative diseases including optic neuropathy, spinal ischemia and stroke, memory and cognition deficits, attention deficit disorder, psychoses including manic disorders, anxiety, depression, hypertension, congestive heart failure, cardiac

ischemia and nasal congestion. See, for example, Hieble et al., "Therapeutic applications of agents interacting with alpha-adrenoceptors, in Alpha-adrenoceptors: molecular biology, biochemistry and pharmacology". *Prog. Basic Clin. Pharmacol.* (Basel, Karger) 8, pp. 180-220(1991). For example,

5 clonidine has been shown to be clinically effective in providing pain relief for postoperative, cancer-associated and neurogenic pain. But, as stated in Maze and Tranquilli, Maze MB and Tranquilli, W. "Alpha-2 adrenoceptor agonists: defining the role in clinical anesthesia". *Anesthesiology* 74, 581-605 (1991), the

10 "full clinical promise" of this and other alpha2 agonists requires the development of compounds that do not cause sedation, hypotension and bradycardia.

The above-listed diseases and disorders are treatable by activation of $\alpha 2B$ or $\alpha 2B/2C$ receptor subtype(s). Therefore, the alpha2 compounds described above that have been shown above not to elicit sedation and

15 cardiovascular effects, are useful and advantageous in the treatment of these conditions.

Amelioration of neuronal degeneration in glaucomatous neuropathy is another example of the novel utility of the compounds of the invention. Recent studies have demonstrated that clonidine and other alpha2 agonists are

20 neuroprotective of retinal cells in several rat models of neuronal degeneration. These models include light-induced photoreceptor degeneration in albino rat, as described in Wen et al, "Alpha2-adrenergic agonists induce basic fibroblast growth factor expression in photoreceptors in vivo and ameliorate light damage." *J. Neurosci.* 16, 5986-5992 and calibrated rat optic nerve injury

25 resulting in secondary loss of retinal ganglion cells, as described in Yoles et al, "Injury-induced secondary degeneration of rat optic nerve can be attenuated

by alpha2-adrenoceptor agonists AGN 191103 and brimonidine". *Invest. Ophthalmol. Vis. Sci.* 37, 540, S114. However, unlike the compounds of the present invention, the doses used in these studies – 0.1 to >1 mg/kg by intraperitoneal or intramuscular injection– also cause sedation and

5 cardiovascular effects. Induction of the expression of basic fibroblast growth factor (bFGF) is considered a sensitive indicator of alpha2 receptor activation in the retina (Wen et al above) and measurement of bFGF induction following topical administration of alpha2 agonists to rat eyes indicates that

10 approximately a 1% dose is necessary to induce a 2-3 fold increase in bFGF levels that correspond with alpha2 agonist mediated neuroprotection (See Wen et al, above, and Lai et al, "Neuroprotective effect of ocular hypotensive agent brimonidine", in *Proceedings of XIth Congress of the European Society of Ophthalmology* (Bologna, Monduzzi Editore), 439-444.) These topical doses of

15 current alpha2 agonists such as clonidine are known to result in systemic side effects such as sedation and hypotension that would prevent their use as ocular neuroprotective agents. Additionally commonly assigned and co-pending application, 08/496,292 filed on 28 June, 1995, discloses and claims the use of certain non-selective $\alpha 2$ -adrenergic agents in treating neural injury, the contents of which are hereby incorporated by reference in their entirety.

20 The compounds of the present invention do not cause sedation and cardiovascular effects following topical administration of doses of at least 3% in monkeys. Thus, neuroprotective concentrations of these compounds can be reached in humans without causing side effects. In fact, as reported below, the compound of Example B-9(b) has been shown to be neuroprotective in the

25 calibrated rat optic nerve injury model of Yoles et al, above. See Table 4, below.

**Table 4: Retinal Ganglion Cell Numbers at 2 Weeks Post-Injury
(cells/microscopic field)**

<u>Control</u> (vehicle i.p.)	<u>Example B-9(b)</u> (0.5 mg/kg i.p.)
33 ± 8	73 ± 12
n = 8	n = 5

- 5 This level of neuroprotection is comparable to the effect seen in previous studies with the standard alpha 2-adrenoceptor agonist, brimonidine, and the neuroprotective agent, MK801.

Example W

- 10 Alleviation of pain including neuropathic pain is another example of a disorder in which the compounds of the invention are useful and advantageous since pain is alleviated without undesirable side effects. Clonidine, an agonist that activates all three alpha2 receptors, has been used clinically for treating chronic pain, but its utility for this indication is limited because it causes sedation and cardiovascular side effects.
- 15 Compounds of the present invention were compared to clonidine and brimonidine in a rodent model of neuropathic pain that is known to be predictive of clinical activity. (See, for example, Kim, S. and Chung, J. "An experimental model for peripheral neuropathy produced by segmental spinal nerve ligation in the rat." *Pain* 50 pp. 355-363 (1992).) Following
- 20 ligation of two spinal nerves, the animals develop a sensitivity to normally non-painful stimuli such as touch. The ability of alpha2 compounds to reverse this sensitivity, called allodynia, was tested 30 minutes after dosing by either intrathecal or intraperitoneal administration. The sedative activity of each compound was also measured using an activity chamber.

The compounds of the invention, exemplified by N-1, are able to alleviate the allodynia without causing sedation, even at very high doses. This is in contrast to clonidine and brimonidine, which cause sedation at doses only slightly higher than their anti-allodynic doses. See tables 5 and 6, below.

Table 5. The anti-allodynic and sedative effects of alpha2-adrenoceptor agonists in rats 30 minutes following intrathecal administration (N=6).

<u>Compound</u>	<u>Dose (ug)</u>	<u>Reversal of Tactile Allodynia (%)</u>	<u>Sedation (%)</u>
<u>Clonidine</u>	0.1	20*	ND
	1	96*	15
	10	ND	60*
<u>N-1</u>	3	13	ND
	30	64*	0
	300	ND	0

* p<0.05 compared to saline control

• ND signifies no data

Table 6. The anti-allodynic and sedative effects of alpha2-adrenoceptor agonists in rats 30 minutes following intraperitoneal administration (N=6).

<u>Compound</u>	<u>Dose (mg/kg)</u>	<u>Reversal of Tactile Allodynia (%)</u>	<u>Sedation (%)</u>
<u>Brimonidine</u>	3	0	ND
	30	37*	24
	300	ND	67*

<u>(Table 6 con't.)</u> <u>Compound</u>	<u>Dose</u> <u>(mg/kg)</u>	<u>Reversal of Tactile</u> <u>Allodynia (%)</u>	<u>Sedation (%)</u>
N-1	3	3	ND
	30	41*	ND
	10,000	ND	0

* p<0.05 compared to saline control

- ND signifies no data


5 The results of these Examples demonstrate that the common side effects of $\alpha 2$ -adrenoceptor drugs are mediated by the $\alpha 2A$ -subtype and that their ocular antihypertensive and other therapeutic actions can be mediated by a subtype other than the $\alpha 2A$ -subtype. Thus, $\alpha 2$ -adrenoceptor compounds of unrelated structural classes, that have in common low functional activity at the $\alpha 2A$ -subtype, lower IOP and elicit other therapeutic actions without dose-limiting side effects.

10 While particular embodiments of the invention have been described, it will be understood, of course, that the invention is not limited thereto since many obvious modifications can be made, and it is intended to include within this invention any such modification as will fall within the scope of the appended claims.

15 Having now described the invention, we claim:

WHAT IS CLAIMED IS:

1. A compound having selective agonist activity at the $\alpha 2B$ or $\alpha 2B/\alpha 2C$ adrenergic receptor subtype(s) as compared to the 2A adrenergic receptor subtype represented by the formula




I

- wherein the dotted lines represent optional double bonds; R is H or lower alkyl; X is S or C(H)R¹, wherein R¹ is H or lower alkyl or R¹ is absent when X is S or when the bond between X and the ring represented by



- is a double bond; Y is O, N, S, (CR¹)_y, wherein y is an integer of from 1 to 3, -CH=CH- or -Y'¹CH₂-, wherein Y'¹ is O, N or S; x is an integer of 1 or 2, wherein x is 1 when R², R³ or R⁴ is bound to an unsaturated carbon atom and x is 2 when R², R³ or R⁴ is bonded to a saturated carbon atom; R² is H, lower alkyl, halogen, hydroxy or lower alkoxy, or, when attached to a saturated carbon atom, R_i may be oxo; R_j and R_k are, each, H, lower alkyl, hydroxy, lower alkoxy, or phenyl or, together, are -(C(R²)_x)_z;


- $-Y^1(C(R^2)_x)z'-;$ $-Y^1(C(R^2)_x)yY^1-;$ $-(C(R^2)_x)-Y^1-(C(R^2)_x)-;$ $-(C(R^2)_x)-Y^1-$
 $(C(R^2)_x)-(C(R^2)_x)-$ and $-Y^1-(C(R^2)_x)-Y^1-(C(R^2)_x)-$ wherein z is an integer of
 from 3 to 5, z' is an integer of from 2 to 4 and x and y are as defined above,
 and further either end of each of these divalent moieties may attach at
 5 either R^3 or R^4 to form the condensed ring structure



- and the ring thus formed may be totally unsaturated, partially unsaturated,
 or totally saturated provided that a ring carbon has no more than 4
 10 valences, nitrogen no more than three and O and S have no more than two,
 and including pharmaceutically acceptable salts thereof.

2. A compound according to claim 1 wherein said compound is
 represented by the formula

15



20

3. A compound according to claim 2 wherein X is $C(H)R^1$.
 4. A compound of claim 3 wherein R^1 is H.

5. A compound of claim 4 wherein R_2 is H and



represents a furanyl radical.

5


6. A compound of claim 5 wherein R^3 and R^4 together are $(CH)_n$.

7. A compound of claim 5 wherein R^3 is H and R^4 is t-butyl.

- 10 8. A compound of claim 5 wherein R^3 and R^4 are H.

9. A compound of claim 5 wherein R^3 is H and R^4 is methyl or ethyl.

- 15 10. A compound of claim 4 wherein R^3 is H and



represents a thienyl radical.

11. A compound of claim 10 wherein R^3 and R^4 , together, represent
20 $(CH)_n$.

12. A compound of claim 10 wherein R^3 is phenyl and R^4 is H.

13. A compound of claim 10 wherein R^3 and R^4 , together, represent $(CH_2)_5S$.

14. A compound of claim 10 wherein R^3 and R^4 are H.

5

15. A compound of claim 10 wherein R^3 and R^4 , together, represent $(CH)_4$.

16. A compound of claim 10 wherein R^3 is H and R^4 is methyl.

10

17. A compound of claim 10 wherein R^3 is bromo and R^4 is H.

18. A compound of claim 4 wherein



15 represents a cyclohexyl radical.

19. A compound of claim 18 wherein R^2 is H, and R^3 and R^4 , together, represent $(CH)_4$.

20. A compound of claim 18 wherein R^2 is H, and R^3 and R^4 , together, represent $(CH)_5S$.

21. A compound of claim 18 wherein R^2 is H, and R^3 and R^4 , together, represent $(CH_2)_4$.

25

22. A compound of claim 18 wherein R^2 is dimethyl, and R^3 and R^4 , together, represent $(CH)_4$.
23. A compound of claim 18 wherein Y is $-CH_2CH(CH_3)-$, R^2 is hydrogen or oxo, and R^3 and R^4 , together, represent $(CH)_4$.
24. A compound of claim 18 wherein R^2 is oxo, and R^3 and R^4 , together, represent $S(CH)_4$.
25. A compound of claim 18 wherein Y is $-CH_2C(CH_3)_2-$, R^2 is hydrogen or oxo, and R^3 and R^4 , together, represent $(CH)_4$.
26. A compound of claim 18 wherein R^2 is oxo, and R^3 and R^4 , together, are $(CH)_4$.
27. A compound of claim 18 wherein R^2 is oxo, and R^3 and R^4 , together, represent $(CH)_2C(OCH_3)CH$.
28. A compound of claim 4 wherein



- represents a cyclopentyl radical.

29. A compound of claim 28 wherein R^2 is H, and R^3 and R^4 , together, represent $(CH)_4$.

30. A compound of claim 28 wherein R^2 is hydrogen, and R^3 and R^4 , together, represent $(CH_2)_3$.

31. A compound of claim 4 wherein




5

represents a benzyl radical.

32. A compound of claim 31 wherein R^2 , R^3 and R^4 are H.

10 33. A compound of claim 1 wherein said compound has the formula



wherein Y is S or O.


34. A compound of claim 33 wherein X is $C(H)R^1$ and R , R^1 , R^2 , R^3 and
15 R^4 are H.

35. A compound of claim 34 wherein Y is O.

36. A compound of claim 35 wherein Y is S.

20

37. A compound of claim 1 wherein said compound has the formula



38. A compound of claim 37 wherein R^3 and R^4 , together, represent
5 $(\text{CH})_x$.
39. A compound of claim 38 wherein Y^1 is O .
40. A compound of claim 39 wherein R^2 is oxo.
- 10 41. A compound of claim 40 wherein X is CH .
42. A compound of claim 40 wherein X is CH_2 .
- 15 43. A compound of claim 39 wherein one of R^2 is hydroxy and the other is H .
44. A compound of claim 39 wherein R^2 is H .
- 20 45. A compound of claim 38 wherein Y^1 is S .

46. A compound of claim 45 wherein X is CH₂.

47. A compound of claim 46 wherein R¹ is oxo.

5 48. A compound of claim 46 wherein R² is H.

49. A compound of claim 45 wherein X is CH and R² is oxo.


50. A compound of claim 3 wherein Y is (CH₂)_n.

10

51. A compound of claim 50 wherein X is CH and R² is oxo.

52. A compound of claim 50 wherein X is CH₂ and R² is H.


15 53. A compound of claim 2 wherein X is S and



is phenyl.

54. A compound of claim 3 wherein R¹ is methyl and

20



is furanyl.


55. A compound of claim 4 wherein Y is $CH_2(CR^1)_2$ wherein R^1 is
5 hydrogen or methyl.

56. A compound of claim 55 wherein R^2 is H.

57. A compound of claim 55 wherein R^2 is oxo.

10

58. A compound of claim 3 wherein R is CH_2 ,




represents a phenyl radical and R^3 and R^4 , together represent $O(CR^1)_2O$.

- 15 59. A compound of claim 2 wherein X is CH_2 ,




represents a cyclopentyl radical and R_1 is oxo.

60. A compound represented by the formula




61. A compound according to claim 1 represented by the formula



- wherein Y is $(R^1)_x$, $R^3 + R^4$ is $(C(R^2))_2$, and X attaches at one of the two positions of the ring indicated by the wavy line with the remaining position being occupied by hydrogen, provided that two double bonds may not occupy the same ring atom.


62. A compound according to claim 61 wherein said compound is represented by the formula




- wherein $(R^2)_x$ is hydrogen or oxo.

63. A compound of claim 61 wherein the structure is


104



64. A compound of claim 62 wherein the structure is




5 65. A compound of claim 2 wherein R is hydrogen, R3 + R4 are –
(C(R²)_x)_n-N-(C(R²)_x)_n-(C(R²)_x)_n-, and X is CHR¹ as represented by the formula



the CHR¹ group attaches at one of the two positions of the ring indicated by the wavy line with the remaining position being occupied by hydrogen, and provided that two double bonds may not occupy the same ring atom.


15

66. A compound of claim 65 wherein said compound has the formula




and (R²)_x is hydrogen or oxo.

67. A compound of claim 65 wherein said compound has the formula




5 and $(R^2)_x$ is hydrogen or oxo.

68. A compound of claim 1 wherein $R_3 + R_4$ are chosen from the group consisting of $-Y^1-(C(R^2)_x)-$, $(C(R^2)_x)-Y^1-$ and $-Y^1(C(R^2)_x)-(C(R^2)_x)-(C(R^2)_x)-$, and Y^1 is N or O or S as represented by the formula




10 wherein X and X' are selected from the group consisting of N, O, and C and
15 at least one of X and X' are N.

69. A compound according to claim 68 wherein said compound is represented by the formula




20 wherein $(R^2)_x$ is hydrogen or oxo.

25 70. A compounds according to claim 68 wherein said compound is represented by the formula



5 wherein $(R^2)_x$ is hydrogen or oxo.

71. A compound having selective agonist activity at the $\alpha 2B$ or $\alpha 2B/2C$ adrenergic receptor or $\alpha 2B$ and $\alpha 2C$ adrenergic receptor subtype(s) as compared to the $\alpha 2A$ adrenergic receptor subtype represented by the



10 formula
and pharmaceutically acceptable salts thereof.

72. A process for administering to a host mammal, including a human being, a pharmaceutical composition containing an effective dose of an active compound to treat or prevent glaucoma without sedating or
15 cardiovascular side effects, wherein said compound has adrenergic activity and is a selective agonist of the $\alpha 2B$ adrenoceptor subtype or $\alpha 2B/\alpha 2C$ adrenoceptor subtype(s) in preference over the $\alpha 2A$ adrenoceptor subtype.

73. A process of claim 72 wherein the active compound has an efficacy
20 relative to a standard full agonist that is at least approximately 0.3 greater at the $\alpha 2B$ or $\alpha 2C$ adrenoceptor subtypes than at the $\alpha 2A$ adrenoceptor subtype and its efficacy at the $\alpha 2A$ adrenoceptor subtype is ≤ 0.4 .

74. A process of claim 72 wherein the active compound is at least ten times more potent at the $\alpha 2B$ or $\alpha 2C$ adrenoceptor subtype than at the $\alpha 2A$ adrenoceptor receptor.

5

75. A process of claim 74 wherein approximately 0.001% to 5% by weight of the active compound is administered topically to the host mammal in daily or twice daily doses.

10 76. A process of claim 75 wherein approximately 0.01% to 3% by weight of the active compound is administered topically to the host mammal in daily or twice daily doses.

15 77. A process of claim 72 wherein said compound has no activity at the $\alpha 2A$ adrenoceptor subtype.

78. A process of claim 72 wherein said compound has no activity at the $\alpha 2A$ and $\alpha 2C$ adrenoceptor subtypes.

20 79. A process for administering to a host mammal, including a human being, a pharmaceutical composition containing an effective dose of an active compound to treat elevated intraocular pressure without sedating or cardiovascular side effects, wherein the compound has adrenergic activity and is a selective agonist of the $\alpha 2B$ or $\alpha 2B/\alpha 2C$ adrenoceptor subtype(s) in
25 preference over the $\alpha 2A$ adrenoceptor receptor subtype.

80. A process of claim 79 wherein the active compound has an efficacy relative to a standard full agonist that is at least approximately 0.3 greater at the $\alpha 2B$ or $2C$ adrenoceptor subtypes than at the $\alpha 2A$ adrenoceptor subtype, and its efficacy at the $\alpha 2A$ adrenoceptor subtype is ≤ 0.4 .

5

81. A process of claim 80 wherein approximately 0.001% to 5% by weight of the active compound is administered topically to the host mammal in daily or twice daily doses.

10 82. A process of claim 81 wherein approximately 0.01% to 3.0% by weight of the active compound is administered topically to the host mammal in daily or twice daily doses.

83. A process of claim 79 wherein said compound has no activity at the
15 $\alpha 2A$ adrenoreceptor subtypes.

84. A process of claim 79 wherein said compound has no activity at the $\alpha 2A$ and $\alpha 2C$ adrenoreceptor subtypes.


20 85. A method of treating a mammal to lower intraocular pressure without having cardiovascular and sedative side effects by selectively agonizing the $\alpha 2B$ adrenoceptor subtype or $\alpha 2B/\alpha 2C$ adrenoceptor subtype(s) in preference to the $\alpha 2A$ adrenoceptor subtype.

25 86. A method of selectively agonizing the $\alpha 2B$ adrenoceptor subtype or $\alpha 2B/\alpha 2C$ adrenoceptor subtypes without agonizing the $\alpha 2A$ adrenoceptor

subtype comprising administering a therapeutically effective amount of a selective $\alpha 2B$ or $\alpha 2B/\alpha 2C$ receptor subtype agonist(s) respectively.

87. An alpha adrenergic agonist that selectively activates the $\alpha 2B$ or $\alpha 2B/\alpha 2C$ receptor subtype(s) in preference to the $\alpha 2A$ receptor subtype.

88. A process according to claims 72, 79, 85 or 91 wherein the active compound is selected from the group consisting of compounds having the formula




wherein the dotted lines represent optional double bonds; R is H or lower alkyl; X is S or C(H)R', wherein R' is H or lower alkyl or R' is absent when X is S or when the bond between X and the ring represented by




is a double bond; Y is O, N, S, (CR¹)_y, wherein y is an integer of from 1 to 3, -CH=CH- or -Y¹CH₂-, wherein Y¹ is O, N or S; x is an integer of 1 or 2, wherein x is 1 when R², R³ or R⁴ is bound to an unsaturated carbon atom and x is 2 when R², R³ or R⁴ is bonded to a saturated carbon atom; R¹ is H, lower alkyl, halogen, hydroxy or lower alkoxy, or, when attached to a

- saturated carbon atom, R_1 may be oxo; R_2 and R_3 are, each, H, lower alkyl, hydroxy, lower alkoxy, or phenyl or, together, are $-(C(R^2)_x)z-$; $-Y^1(C(R^2)_x)z'-$; $-Y^1(C(R^2)_x)Y^1-$; $-(C(R^2)_x)-Y^1-(C(R^2)_x)-$; $-(C(R^2)_x)-Y^1-(C(R^2)_x)-(C(R^2)_x)-$ and $-Y^1-(C(R^2)_x)-Y^1-(C(R^2)_x)-$ wherein z is an integer of
- 5 from 3 to 5, z' is an integer of from 2 to 4 and x and y are as defined above, and further either end of each of these divalent moieties may attach at either R^3 or R^4 to form the condensed ring structure




- and the ring thus formed may be totally unsaturated, partially unsaturated,
- 10 or totally saturated provided that a ring carbon has no more than 4 valences, nitrogen no more than three and O and S have no more than two;
- or




- 15 wherein W is a bicyclic radical selected from the group consisting of

111



wherein R^5 , R^6 , R^7 and R^8 are selected from the group consisting of H and lower alkyl provided that at least one of R^5 and R^6 or R^6 and R^7 are $OC(R^9)C(R^9)N(R)$ to form a condensed ring with



5

wherein R^9 is H, lower alkyl or oxo
and



wherein R^{10} is H, lower alkyl, phenyl or lower alkyl substituted phenyl,
10 and Z is O or NH.

89. A process for administering to a host mammal, including a human being, a pharmaceutical composition containing an effective dose of an active compound, having adrenergic activity, to treat or prevent glaucoma
15 wherein the active compound has the biological property that the compound is a selective agonist of $\alpha 2B$ or $\alpha 2B/\alpha 2C$ receptor subtype(s) in

preference over the $\alpha 2A$ receptor subtype, said selectivity being measured in an assay using cells that naturally express the individual $\alpha 2$ subtypes or have had one of the subtypes introduced, the receptors being human or from a species that has been shown to have a similar pharmacology, and in
5 which assay the efficacy relative to a standard compound of the active compound at the $\alpha 2B$ or $\alpha 2C$ receptor subtype is measured to be at least 0.3 greater than the efficacy relative to the standard compound of the active compound at the $\alpha 2A$ receptor subtype and its efficacy at the $\alpha 2A$ receptor subtype is ≤ 0.4 , and/or the active compound is at least approximately 10
10 times more potent at the $\alpha 2B$ or $\alpha 2C$ receptor subtypes than at the $\alpha 2A$ receptor subtype.

90. A process of claim 89 wherein approximately 0.001% to 5% by weight of the active compound is administered topically to the host
15 mammal per day.

91. A process for administering to a host mammal, including a human being, a pharmaceutical composition containing an effective dose of an active compound to treat or prevent muscle spasticity including
20 hyperactive micturition, diarrhea, diuresis, withdrawal syndromes, pain including neuropathic pain, neurodegenerative diseases including optic neuropathy, spinal ischemia and stroke, memory and cognition deficits, attention deficit disorder, psychoses including manic disorders, anxiety, depression, hypertension, congestive heart failure, cardiac ischemia and
25 nasal congestion without sedating or cardiovascular side effects, wherein said compound has adrenergic activity and is a selective agonist of the $\alpha 2B$

or $\alpha 2B$ / $\alpha 2C$ adrenoceptor receptor subtype(s) in preference over the $\alpha 2A$ adrenoceptor receptor subtype.

92. A process for administering to a host mammal, including a human
5 being, a pharmaceutical composition containing an effective dose of an
active compound, having adrenergic activity, to treat muscle spasticity
including hyperactive micturition, diarrh ea, diuresis, withdrawal
syndromes, pain including neuropathic pain, neurodegenerative diseases
including optic neuropathy, spinal ischemia and stroke, memory and
10 cognition deficits, attention deficit disorder, psychoses including manic
disorders, anxiety, depression, hypertension, congestive heart failure,
cardiac ischemia and nasal congestion without sedating or cardiovascular
side effects wherein the active compound has the biological property that
the compound is a selective agonist of $\alpha 2B$ or $\alpha 2B$ / $\alpha 2C$ receptor subtype(s)
15 in preference over the $\alpha 2A$ receptor subtype, said selectivity being
measured in an assay using cells that naturally express the individual $\alpha 2$
subtypes or have had one of the subtypes introduced, the receptors being
human or from a species that has been shown to have a similar
pharmacology, and in which assay the efficacy relative to a standard
20 compound of the active compound at the $\alpha 2B$ or the $\alpha 2C$ receptor subtype
is measured to be at least 0.3 greater than the efficacy relative to the
standard compound of the active compound at the $\alpha 2A$ receptor subtype
and its efficacy at the $\alpha 2A$ receptor is ≤ 0.4 , and/or the active compound is
at least approximately 10 times more potent at the $\alpha 2B$ or $\alpha 2C$ receptor
25 subtypes than at the $\alpha 2A$ receptor subtype.

93. A process for administering to a host mammal, including a human being, a pharmaceutical composition containing an effective dose of an active compound, having adrenergic activity, to treat muscle spasticity including hyperactive micturition, diarrhea, diuresis, withdrawal syndromes, pain including neuropathic pain, neurodegenerative diseases including optic neuropathy, spinal ischemia and stroke, memory and cognition deficits, attention deficit disorder, psychoses including manic disorders, anxiety, depression, hypertension, congestive heart failure, cardiac ischemia and nasal congestion without sedating or cardiovascular side effects wherein the active compound has the biological property that the compound is a selective agonist of $\alpha 2B$ or $\alpha 2B / \alpha 2C$ receptor subtype(s) in preference over the $\alpha 2A$ receptor subtype, said selectivity being measured in an RSAT assay in which activation of the $\alpha 2A$ and $\alpha 2C$ receptor subtype by the test compound is compared to brimonidine and the $\alpha 2B$ receptor subtype is compared to oxymetazoline and wherein the respective $\alpha 2A$, $\alpha 2B$ and $\alpha 2C$ receptor subtypes are expressed in NIH-3T3 cells, and in which assay the efficacy relative to brimonidine of the active compound at the $\alpha 2C$ receptor subtype or the efficacy relative to oxymetazoline of the active compound at the $\alpha 2B$ receptor subtype is measured to be at least 0.3 greater than the efficacy relative to brimonidine of the active compound at the $\alpha 2A$ receptor subtype, and its efficacy at the $\alpha 2A$ receptor subtype is ≤ 0.4 and/or the active compound is at least approximately 10 times more potent at the $\alpha 2B$ or $\alpha 2C$ receptor subtypes than at the $\alpha 2A$ receptor subtype.

94. A process of claim 74 wherein the active compound is at least one hundred times more potent at the $\alpha 2B$ or $\alpha 2C$ adrenoceptor subtype than at the $\alpha 2A$ adrenoceptor subtype.

INTERNATIONAL SEARCH REPORT

Int. l. Application No.
PCT/US 98/25669

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07D233/56 A61K31/415 C07D401/06 C07D403/06 C07D409/06 C07D405/06 C07D233/84 C07D413/12 C07D495/04 //(C07D495/04, 335:00, 333:00)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 07866 A (TOKYO TANABE CO MPANY LIMITED) 14 April 1994 see abstract	1-4, 18-27, 37
X	WO 97 35858 A (ORTHO PHARMA CORP) 2 October 1997 see page 7; claims	1-4, 10-17, 72
X	US 5 621 113 A (BOYD ROBERT E ET AL) 15 April 1997 see column 6; claims	1-4, 10-17, 72
X	WO 97 12874 A (ORION-YHTYMÄ) 10 April 1997 see page 13 - page 15, line 2; claims	1-4, 18-30, 72
-/-		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "Z" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
29 March 1999		12/04/1999
Name and mailing address of the ISA European Patent Office, P.B. 5618 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3018		Authorized officer Henry, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/25669

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 304 910 A (SEARLE & CO) 1 March 1989 see page 2, line 50 - page 4, line 20; claims	1-4,72
A	EP 0 194 984 A (CONTINENTAL PHARMA) 17 September 1986 see page 1 - page 2, line 15; claims	1-4,72
A	CHEMICAL ABSTRACTS, vol. 112, no. 15, 9 April 1990 Columbus, Ohio, US; abstract no. 139033n, page 721; XP002098179 see abstract & JP 01 242571 A (MITSUI PETROCHEMICAL INDUSTRIES LTD.) 27 September 1989	1-4,72

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 25669

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 85,86
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 85,86
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/25669

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9407866 A	14-04-1994	AU 5118493 A JP 6192233 A	26-04-1994 12-07-1994
WO 9735858 A	02-10-1997	US 5750720 A AU 2422997 A NO 984481 A	12-05-1998 17-10-1997 23-11-1998
US 5621113 A	15-04-1997	AU 2544097 A EP 0889891 A NO 984482 A WO 9735857 A	17-10-1997 13-01-1999 24-11-1998 02-10-1997
WO 9712874 A	10-04-1997	AU 7132796 A CA 2231535 A CN 1198741 A CZ 9801018 A EP 0888309 A LT 98057 A, B LV 12108 A NO 981496 A PL 328167 A SI 9620111 A	28-04-1997 10-04-1997 11-11-1998 16-09-1998 07-01-1999 25-09-1998 20-08-1998 02-04-1998 18-01-1999 31-10-1998
EP 0304910 A	01-03-1989	US 4882343 A JP 1071871 A	21-11-1989 16-03-1989
EP 0194984 A	17-09-1986	LU 85747 A AT 51618 T AU 597239 B AU 5269686 A CA 1273942 A DK 39286 A FI 860369 A GR 860223 A JP 61176571 A PT 81910 B US 4738979 A	04-08-1986 15-04-1990 31-05-1990 31-07-1986 11-09-1990 29-07-1986 29-07-1986 24-04-1986 08-08-1986 13-05-1987 19-04-1988

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.